arXiv:cond-mat/9703079v2 [cond-mat.supr-con] 6 Feb 2002

The Many-Electron System in the Forward, Exchange and BCS Approximation

by

Detlef Lehmann*
University of British Columbia
Department of Mathematics
Vancouver, B.C.
V6T 1Z2, Canada

Abstract: The nonrelativistic many-electron system in the forward, exchange and BCS approximation is considered. In this approximation, the model is explicitly solvable for arbitrary space dimension d. The partition function and the correlation functions are given by finite-dimensional integral representations. Renormalization effects as well as symmetry breaking can be seen explicitly. It is shown that the usual mean field approach, based on approximating the Hamiltonian by a quadratic expression, may be misleading if the electron-electron interaction contains higher angular momentum terms and the space dimension is $d = 3$. The perturbation theory of the solvable model is discussed. There are cases where the logarithm of the partition function has positive radius of convergence but the sum of all connected diagrams has radius of convergence zero implying that the linked cluster theorem is not applicable in these cases.

---

* present address: TU Berlin, FB Mathematik Ma 7-2, Strasse des 17. Juni 136, D-10623 Berlin; e-mail: lehmann@math.tu-berlin.de

I. Introduction

In this paper we consider the nonrelativistic many-electron system in the forward, exchange and BCS approximation. In this approximation, which is still quartic in the annihilation and creation operators, the model can be solved explicitly. The partition function and the correlation functions are given by finite-dimensional integral representations. We work in the quantum grand canonical ensemble and start with positiv temperature $T = \frac{1}{\beta} > 0$ and finite volume $L^d < \infty$.

The standard model of the many-electron system in d space dimensions is given by the Hamiltonian

$$H = H_0 + H_{\rm int} \tag{I.1}$$

where

$$H_0 = \tfrac{1}{L^d} \sum_{\mathbf{k},\sigma} e_{\mathbf{k}}\, a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma} \tag{I.2}$$

and

$$H_{\rm int} = \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p},\mathbf{q}} U(\mathbf{k}-\mathbf{p})\, a^+_{\mathbf{k}\sigma} a_{\mathbf{p}\sigma} a^+_{\mathbf{q}-\mathbf{k}\tau} a_{\mathbf{q}-\mathbf{p}\tau} \tag{I.3}$$

We assume U to be short range, that is U is L^1 in coordinate space. The energy momentum relation $e_{\mathbf{k}}$ is given by

$$e_{\mathbf{k}} = \tfrac{\mathbf{k}^2}{2m} - \mu \tag{I.4}$$

or may be substituted by a more general expression which satisfies $e_{\mathbf{k}} = e_{-\mathbf{k}}$. The parameter $\mu > 0$, the chemical potential, is present since we are working in the grand canonical ensemble and is determined by the density of the system. Since we are in finite volume, the spatial momenta $\mathbf{k}$ range over some subset of $\left(\frac{2\pi}{L}\mathbf{Z}\right)^d$. The physics of the nonrelativistic many-electron system is determined by momenta close to the Fermi surface

$$F = \left\{\, \mathbf{k} \in \left(\tfrac{2\pi}{L}\mathbf{Z}\right)^d \;\middle|\; e_{\mathbf{k}} = 0 \,\right\} \tag{I.5}$$

so we impose a fixed ultraviolet cuttoff and choose

$$\mathbf{k} \in M_\omega = \left\{\, \mathbf{k} \in \left(\tfrac{2\pi}{L}\mathbf{Z}\right)^d \;\middle|\; |e_{\mathbf{k}}| \le \omega \,\right\} \tag{I.6}$$

In the context of conventional superconductivity the cuttoff ω is referred to as the Debye frequency.

The normalizations choosen in the definition of the creation and annihilation operators are such that the anticommutation relations read

$$\{a_{\mathbf{k}\sigma}, a^+_{\mathbf{k}'\sigma'}\} = L^d \delta_{\mathbf{k},\mathbf{k}'} \delta_{\sigma,\sigma'} \tag{I.7}$$

In particular, if $\Omega_F = \prod_{\substack{\mathbf{k}\sigma \\ e_{\mathbf{k}}<0}} a^+_{\mathbf{k}\sigma}\,\mathbf{1}$ is the zero temperature ground state of the noninteracting system, then $(\Omega_F, a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}\Omega_F) = L^d\theta(-e_{\mathbf{k}})$ such that $(\Omega_F, H_0\Omega_F) = \sum_{\mathbf{k},\sigma} e_{\mathbf{k}}\,\theta(-e_{\mathbf{k}}) \approx L^d \int \frac{d^d\mathbf{k}}{(2\pi)^d}\, e_{\mathbf{k}}\,\theta(-e_{\mathbf{k}})$ is indeed an extensive quantity as it should be. Here $\theta(\varepsilon)$ is the step function being one for $\varepsilon > 0$ and zero otherwise.

We are interested in the grand canonical partition function

$$Z = Z(\beta, L, U) = Tr\, e^{-\beta H} \tag{I.8}$$

which may be normalized by $Tr\, e^{-\beta H_0}$ such that $Z(U = 0) = 1$, and in particular in the two point function

$$\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}'\sigma'}\rangle_{\beta,L} = \frac{Tr\, e^{-\beta H} a^+_{\mathbf{k}\sigma} a_{\mathbf{k}'\sigma'}}{Tr\, e^{-\beta H}} \tag{I.9}$$

which gives the momentum distribution of the system. Recall that in the free ($U = 0$) system, the ideal Fermi gas,

$$\lim_{\beta\to\infty} \langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}'\sigma'}\rangle_{\beta,L} = L^d \delta_{\mathbf{k},\mathbf{k}'}\delta_{\sigma,\sigma'}\theta(-e_{\mathbf{k}}) \tag{I.10}$$

Since we are in the quantum grand canonical ensemble, the traces in (I.8,9) are to be taken over the Fock space $\mathcal{F} = \oplus_{n=0}^{\infty}\mathcal{F}_n$ where

$$\mathcal{F}_n = \left\{ F_n \in L^2\left(\left[[0,L]^d \times \{\uparrow,\downarrow\}\right]^n\right) \,\middle|\, F_n(\mathbf{x}_{\pi 1}\sigma_{\pi 1}, \cdots, \mathbf{x}_{\pi n}\sigma_{\pi n}) = \mathrm{sgn}\pi\; F_n(\mathbf{x}_1\sigma_1, \cdots, \mathbf{x}_n\sigma_n)\right\}$$

The fact that the physical system has a fixed number of particles N is expressed by requireing that the expectation value $\langle \mathbf{N}\rangle_{\beta,L}$ of the number operator $\mathbf{N} = \frac{1}{L^d}\sum_{\mathbf{k}\sigma} a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}$ (which is extensive, see above) is equal to the number of particles, $\langle \mathbf{N}\rangle_{\beta,L} = N$, which determines μ as a function of the density $\frac{N}{L^d}$.

As usual, the quartic part $H_{\rm int}$ (I.3) of the Hamiltonian H (I.1) may be represented by the following four legged diagram:

$H_{\rm int}$:


$\mathbf{k}, \sigma$
$\mathbf{q}-\mathbf{k}, \tau$
$U(\mathbf{k}-\mathbf{p})$
$\mathbf{p}, \sigma$
$\mathbf{q}-\mathbf{p}, \tau$

Because of conservation of momentum, there are three independent momenta here labelled with $\mathbf{k}, \mathbf{p}$ and $\mathbf{q}$. Then one can consider the following three limiting cases with only two independ momenta:



That is, one may consider the approximation

$$H_{\rm int} \approx H_{\rm forw} + H_{\rm ex} + H_{\rm BCS} \tag{I.11}$$

where

$$\begin{aligned} H_{\rm forw} &= \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p},\mathbf{q}} U(\mathbf{k}-\mathbf{p})\, \delta_{\mathbf{k},\mathbf{p}}\, a^+_{\mathbf{k}\sigma} a_{\mathbf{p}\sigma} a^+_{\mathbf{q}-\mathbf{k}\tau} a_{\mathbf{q}-\mathbf{p}\tau} \\ &= \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p}} U(\mathbf{0})\, a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma} a^+_{\mathbf{p}\tau} a_{\mathbf{p}\tau} \end{aligned} \tag{I.12}$$

$$\begin{aligned} H_{\rm ex} &= \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p},\mathbf{q}} U(\mathbf{k}-\mathbf{p})\, \delta_{\mathbf{k},\mathbf{q}-\mathbf{p}}\, a^+_{\mathbf{k}\sigma} a_{\mathbf{p}\sigma} a^+_{\mathbf{q}-\mathbf{k}\tau} a_{\mathbf{q}-\mathbf{p}\tau} \\ &= \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p}} U(\mathbf{k}-\mathbf{p})\, a^+_{\mathbf{k}\sigma} a_{\mathbf{p}\sigma} a^+_{\mathbf{p}\tau} a_{\mathbf{k}\tau} \end{aligned} \tag{I.13}$$

$$\begin{aligned} H_{\rm BCS} &= \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p},\mathbf{q}} U(\mathbf{k}-\mathbf{p})\, \delta_{\mathbf{q},\mathbf{0}}\, a^+_{\mathbf{k}\sigma} a_{\mathbf{p}\sigma} a^+_{\mathbf{q}-\mathbf{k}\tau} a_{\mathbf{q}-\mathbf{p}\tau} \\ &= \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p}} U(\mathbf{k}-\mathbf{p})\, a^+_{\mathbf{k}\sigma} a_{\mathbf{p}\sigma} a^+_{-\mathbf{k}\tau} a_{-\mathbf{p}\tau} \end{aligned} \tag{I.14}$$

Let us shortly make a comment on the volume factors. In (I.12-14), we only introduced some Kroenecker delta's but we did not cancel a volume factor L^d. That this is the right thing to do, that is, that the left hand side as well as the right hand side of (I.11) is indeed proportional to the volume may be seen in the easiest way for the forward term. On a fixed n particle space $\mathcal{F}_n$ the interacting part $H_{\rm int}$ is a multiplication operator given by

$$H_{{\rm int}|_{\mathcal{F}_n}} = \tfrac{1}{2} \sum_{\substack{i,j=1 \\ i\neq j}}^{n} U(\mathbf{x}_i - \mathbf{x}_j) \tag{I.15}$$

Let $\delta_{\mathbf{y}}(\mathbf{x}) = \delta(\mathbf{x}-\mathbf{y})$. Then $\varphi(\mathbf{x}_1, \cdots, \mathbf{x}_n) = \delta_{\mathbf{y}_1} \wedge \cdots \wedge \delta_{\mathbf{y}_n}(\mathbf{x}_1, \cdots, \mathbf{x}_n)$ is an eigenfunction of $H_{\rm int}$ with eigenvalue

$$E = \tfrac{1}{2} \sum_{\substack{i,j=1 \\ i\neq j}}^{n} U(\mathbf{y}_i - \mathbf{y}_j) = \tfrac{1}{2L^d} \sum_{\substack{i,j=1 \\ i\neq j}}^{n} \sum_{\mathbf{q}} e^{i(\mathbf{y}_i - \mathbf{y}_j)\mathbf{q}}\, U(\mathbf{q}) \tag{I.16}$$

which is, for $U \in L^1$, proportional to n or to the volume L^d for constant density. One finds that φ is also an eigenvector of the forward term, $H_{\rm forw}\varphi = E_{\rm forw}\varphi$ where $E_{\rm forw}$ is obtained from (I.16) by putting $\mathbf{q} = 0$ without cancelling a volume factor,

$$E_{\rm forw} = \tfrac{1}{2L^d} \sum_{\substack{i,j=1 \\ i\neq j}}^{n} U(\mathbf{q}=0) \tag{I.17}$$

which is also proportional to the volume.

We now come to the exact definition of the model which is solved in this paper. To do so, we need the functional integral representation of the perturbation series for the partition function. It is summarized in the following theorem which is fairly standard. One may look in [FKT1] for a nice and clean proof.

Theorem I.1: *Let $H = H_0 + H_{\rm int}$ be the Hamiltonian (I.1-3), let*

$$Z = Z(\beta, L) = Tr\, e^{-\beta(H_0 + H_{\rm int})} / Tr\, e^{-\beta H_0} \tag{I.18}$$

be the normalized grand canonical partition function and let

$$\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma} \rangle_{\beta,L} = \frac{Tr\, e^{-\beta H} a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}}{Tr\, e^{-\beta H}} \tag{I.19}$$

a) *The partition function Z has the following perturbation series*

$$Z = \sum_{n=0}^{\infty} \frac{(-\frac{1}{\beta L^d})^n}{n!} \sum_{\sigma_1 \cdots \sigma_{2n}} \sum_{\substack{k_1 \cdots k_{2n} \\ p_1 \cdots p_{2n}}} \prod_{i=1}^{n} \left\{ \delta_{k_{2i-1}+k_{2i}, p_{2i-1}+p_{2i}} U(\mathbf{k}_{2i} - \mathbf{p}_{2i}) \right\} \times$$

$$\det\left[\delta_{\sigma_i,\sigma_j} \delta_{k_i,p_j}\, C(k_i)\right]_{1\le i,j\le 2n} \tag{I.20}$$

Here $k = (k_0, \mathbf{k}) \in \frac{\pi}{\beta}(2\mathbf{Z}+1) \times \left\{ \mathbf{k} \in \left(\frac{2\pi}{L}\mathbf{Z}\right)^d \,\middle|\, |e_{\mathbf{k}}| \le \omega \right\}$ and the covariance C is given by

$$C(k_0, \mathbf{k}) = \frac{1}{ik_0 - e_{\mathbf{k}}} \tag{I.21}$$

If $\|U(\mathbf{x})\|_{L^1} \le c\,(\beta L^d)^{-1}$ or $\|U(\mathbf{x})\|_{L^\infty} \le c\,(\beta L^d)^{-2}$, c a constant, then (I.20) converges.

b) *Let*

$$d\mu_C(\psi,\bar\psi) = \prod_{k,\sigma} \tfrac{\beta L^d}{ik_0-e_{\mathbf{k}}}\; e^{-\frac{1}{\beta L^d}\sum_{k,\sigma}(ik_0-e_{\mathbf{k}})\bar\psi_{k,\sigma}\psi_{k,\sigma}} \prod_{k,\sigma} d\psi_{k,\sigma}d\bar\psi_{k,\sigma} \tag{I.22}$$

be the Grassmann Gaussian measure with covariance C. Then the perturbation series (I.20) can be rewritten as

$$Z = \int e^{-\mathcal{V}_{\rm int}(\psi,\bar\psi)} d\mu_C(\psi,\bar\psi) \tag{I.23}$$

where

$$\mathcal{V}_{\rm int}(\psi,\bar\psi) = \tfrac{1}{(\beta L^d)^3} \sum_{\sigma_1\sigma_2} \sum_{\substack{k_1,k_2\\ p_1,p_2}} \delta_{k_1+k_2,p_1+p_2} U(\mathbf{k}_2-\mathbf{p}_2) \bar\psi_{k_1\sigma_1}\bar\psi_{k_2\sigma_2}\psi_{p_1\sigma_1}\psi_{p_2\sigma_2} \tag{I.24}$$

c) *The momentum distribution (I.19) at temperature $T=\frac{1}{\beta}$ is given by*

$$\langle a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}\rangle_{\beta,L} = \lim_{\substack{\epsilon\to 0\\ \epsilon<0}} \tfrac{1}{\beta} \sum_{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)} \langle \bar\psi_{k_0\mathbf{k}\sigma}\psi_{k_0\mathbf{k}\sigma}\rangle_{\beta,L,\epsilon} \tag{I.25}$$

where

$$\langle \bar\psi_{k\sigma}\psi_{k\sigma}\rangle_{\beta,L,\epsilon} = \tfrac{1}{Z_\epsilon} \int \bar\psi_{k\sigma}\psi_{k\sigma}\; e^{-\mathcal{V}_{\rm int}(\psi,\bar\psi)} d\mu_{C_\epsilon}(\psi,\bar\psi) \tag{I.26}$$

Here $C_\epsilon(k) = \frac{e^{-i\epsilon k_0}}{ik_0-e_{\mathbf{k}}}$ and $Z_\epsilon = \int e^{-\mathcal{V}_{\rm int}} d\mu_{C_\epsilon}$.

Remarks: **(i)** A bound from which convergence of the perturbation series follows for sufficiently small U is easiest obtained in coordinate space. There the perturbation series reads

$$\begin{aligned} Z &= \int e^{-\int d\xi_1 d\xi_2 (\bar\psi\psi)(\xi_1)U(\xi_1-\xi_2)(\bar\psi\psi)(\xi_2)} d\mu_C(\psi,\bar\psi) \\ &= \sum_{n=0}^{\infty} \tfrac{1}{n!} \int d\xi_1\cdots d\xi_{2n} \prod_{i=1}^{n} U(\xi_{2i-1}-\xi_{2i})\; \det\left[C(\xi_i,\xi_j)\right]_{1\le i,j\le 2n} \end{aligned} \tag{I.27}$$

where $\xi=(x_0,\mathbf{x},\sigma)$, $\int d\xi = \sum_{\sigma\in\{\uparrow,\downarrow\}}\int_{[0,\beta]}dx_0\int_{[0,L]^d}d^d\mathbf{x}$, $U(\xi)=U(x)=\delta(x_0)U(\mathbf{x})$, $C(\xi,\xi')=\delta_{\sigma,\sigma'}C(x-x')$ where

$$C(x_0,\mathbf{x}) = \tfrac{1}{L^d} \sum_{\mathbf{k}\in M_\omega} e^{i\mathbf{k}\mathbf{x}} e^{-e_{\mathbf{k}}x_0}\left[\theta(-x_0)\theta_\beta(-e_{\mathbf{k}}) - \theta(x_0)\theta_\beta(e_{\mathbf{k}})\right] \tag{I.28}$$

and θ_β is an approximate step function given by $\theta_\beta(\varepsilon)=\frac{1}{1+e^{-\beta\varepsilon}}$. First observe that, since $\sup_{x_0\in[0,\beta]}\left|e^{-e_{\mathbf{k}}x_0}[\theta(-x_0)\theta_\beta(-e_{\mathbf{k}})-\theta(x_0)\theta_\beta(e_{\mathbf{k}})]\right| \le 1$ for all $\mathbf{k}$, one has $\|C\|_\infty \le \frac{1}{L^d}\sum_{\mathbf{k}\in M_\omega} 1 < \infty$ if we choose a fixed ultraviolet cuttoff in (I.6). The determinant can be bounded using $|\det[\vec{a}_1 \cdots \vec{a}_r]| \le \Pi_{i=1}^r \|\vec{a}_i\|_2$ where the $\vec{a}_i$ are r component vectors. That is

$$\left|\det\left[C(\xi_i,\xi_j)\right]_{1\le i,j\le 2n}\right| \le \prod_{i=1}^{2n}\Big\{\sum_{j=1}^{2n}|C(\xi_i,\xi_j)|^2\Big\}^{\frac{1}{2}} \le \prod_{i=1}^{2n}(2n\|C\|_\infty^2)^{\frac{1}{2}} = (2n\|C\|_\infty^2)^n \tag{I.29}$$

which gives

$$|Z| \le \sum_{n=0}^{\infty}\left(\tfrac{e}{n}\right)^n(4\beta L^d)^n\|U\|_{L^1}^n(2n\|C\|_\infty^2)^n = \sum_{n=0}^{\infty}\left(8e\|C\|_\infty^2\beta L^d\right)^n\|U\|_{L^1}^n \tag{I.30}$$

(ii) The ϵ-limit shows up in (I.25) because of

$$\begin{aligned} C(x_0,\mathbf{x}) &\overset{x_0\ne 0}{=} \tfrac{1}{L^d}\sum_{\mathbf{k}} e^{i\mathbf{kx}}e^{-e_{\mathbf{k}}x_0}\left[\theta_\beta(-e_{\mathbf{k}})\theta(-x_0)-\theta_\beta(e_{\mathbf{k}})\theta(x_0)\right] \\ &= \tfrac{1}{\beta L^d}\sum_{\mathbf{k},k_0} e^{i\mathbf{kx}-ik_0x_0}\tfrac{1}{ik_0-e_{\mathbf{k}}} \end{aligned} \tag{I.31}$$

but

$$C(0,\mathbf{x}) = \tfrac{1}{L^d}\sum_{\mathbf{k}} e^{i\mathbf{kx}}\theta_\beta(-e_{\mathbf{k}}) = \lim_{\substack{\epsilon\to 0\\ \epsilon<0}}\tfrac{1}{\beta L^d}\sum_{\mathbf{k},k_0} e^{i\mathbf{kx}-ik_0\epsilon}\tfrac{1}{ik_0-e_{\mathbf{k}}}\ , \tag{I.32}$$

see the proof of Lemma A1 in the appendix for a more detailed discussion.

(iii) Whereas the momenta in the Hamiltonian are d dimensional, the variables in the perturbation series are $d+1$ dimensional. The additional k_0 variables in (I.20) are the Fourier transform of the x_0 variables in (I.27) which in turn enter the perturbation series because of

$$\tfrac{d}{d\lambda}{}_{|0}\, e^{-\beta(H_0+\lambda H_{\rm int})} = \int_0^\beta e^{-x_0H_0}H_{\rm int}\,e^{-(\beta-x_0)H_0}dx_0\ , \tag{I.33}$$

that is, because $H_{\rm int}$ does not commute with H_0.

As remarked under (iii), the momenta in

$$H_{\rm int} = \tfrac{1}{L^{3d}}\sum_{\sigma,\tau\in\{\uparrow,\downarrow\}}\sum_{\mathbf{k},\mathbf{p},\mathbf{q}} U(\mathbf{k}-\mathbf{p})\, a^+_{\mathbf{k}\sigma}a_{\mathbf{p}\sigma}a^+_{\mathbf{q}-\mathbf{k}\tau}a_{\mathbf{q}-\mathbf{p}\tau} \tag{I.3}$$

are d dimensional, but the variables in

$$\mathcal{V}_{\rm int} = \tfrac{1}{(\beta L^d)^3}\sum_{\sigma,\tau\in\{\uparrow,\downarrow\}}\sum_{k,p,q} U(\mathbf{k}-\mathbf{p})\,\bar\psi_{k\sigma}\psi_{p\sigma}\bar\psi_{q-k\tau}\psi_{q-p\tau} \tag{I.34}$$

are $d+1$ dimensional. On the Hamiltonian level, the forward, exchange and BCS approximation (I.11) does not lead to an explicitly solvable model, but on the functional integral level it does. This is the main result of this paper.

Theorem I.2: *Let*

$$\{\mathcal{V}_{\rm forw}+\mathcal{V}_{\rm ex}+\mathcal{V}_{\rm BCS}\}(\psi,\bar\psi)=\tfrac{1}{(\beta L^d)^3}\sum_{\sigma\tau}\sum_{kpq}U(\mathbf{k}-\mathbf{p})\,[\delta_{k,p}+\delta_{k,q-p}+\delta_{q,0}]\,\bar\psi_{k\sigma}\psi_{p\sigma}\bar\psi_{q-k\tau}\psi_{q-p\tau} \tag{I.35}$$

Then the approximation

$$\mathcal{V}_{\rm int}\approx\mathcal{V}_{\rm forw}+\mathcal{V}_{\rm ex}+\mathcal{V}_{\rm BCS} \tag{I.36}$$

in (I.23,26) makes the model explicitly solvable. The partition function, the two point functions and the generating functional for the connected amputated Greens functions are given by finite-dimensional integral representations. The dimension of these representations is proportional to J if the electron electron interaction is given by

$$U(\mathbf{k}-\mathbf{p})=\begin{cases}\dfrac{1}{2}\displaystyle\sum_{\ell=-j}^{j}\lambda_{|\ell|}e^{i\ell\varphi_{\mathbf{k}}}e^{-i\ell\varphi_{\mathbf{p}}}+\dfrac{\lambda_0}{2} & \text{if } d=2\\ \displaystyle\sum_{\ell=0}^{j}\sum_{m=-\ell}^{\ell}\lambda_\ell\bar Y_{\ell m}(\mathbf{k}')\,Y_{\ell m}(\mathbf{p}') & \text{if } d=3\end{cases}\quad=:\ \sum_{l=0}^{J}\lambda_l\,y_l(\mathbf{k})\,\bar y_l(\mathbf{p}) \tag{I.37}$$

Here $\mathbf{k}=|\mathbf{k}|(\cos\varphi_{\mathbf{k}},\sin\varphi_{\mathbf{k}})$ *for* $d=2$ *and* $\mathbf{k}'=\mathbf{k}/|\mathbf{k}|$ *for* $d=3$.

The general solution is written down in Theorem II.1. In the following, we summarize the results for $\mathcal{V}_{\rm int}\approx\mathcal{V}_{\rm BCS}$ and for an electron-electron interaction given by (I.37). This case is treated in section III.1.

In that case the two point function (I.26) becomes (we suppress the ϵ):

$$\langle\bar\psi_{p\uparrow}\psi_{p'\uparrow}\rangle_{\beta,L}=\beta L^d\delta_{p,p'}\frac{\int\ F_{\mathbf{p}}^{\uparrow\uparrow}(\phi)\,e^{-\beta L^dV(\phi)}\prod\limits_{\sigma\tau}\prod\limits_{l=0}^{J}du^l_{\sigma\tau}dv^l_{\sigma\tau}}{\int e^{-\beta L^dV(\phi)}\prod\limits_{\sigma\tau}\prod\limits_{l=0}^{J}du^l_{\sigma\tau}dv^l_{\sigma\tau}} \tag{I.38}$$

Here $\phi^l_{\sigma\tau}=u^l_{\sigma\tau}+iv^l_{\sigma\tau}$, $\sigma\tau\in\{\uparrow\uparrow,\downarrow\downarrow,\uparrow\downarrow\}$,

$$F_{\mathbf{p}}^{\uparrow\uparrow}(\phi)=\tfrac{(ip_0+e_{\mathbf{p}})[p_0^2+e_{\mathbf{p}}^2+\Phi_{-\mathbf{p}\uparrow\downarrow}\bar\Phi_{-\mathbf{p}\uparrow\downarrow}+\Phi_{\mathbf{p}\downarrow\downarrow}\bar\Phi_{\mathbf{p}\downarrow\downarrow}]}{(p_0^2+e_{\mathbf{p}}^2+\Omega^+_{\mathbf{p}})(p_0^2+e_{\mathbf{p}}^2+\Omega^-_{\mathbf{p}})} \tag{I.39}$$

$$\Phi_{\mathbf{k}\uparrow\downarrow}=\sum_{l=0}^{J}\lambda_l^{\frac12}\phi^l_{\uparrow\downarrow}\,y_l(\mathbf{k}')\,,\qquad\bar\Phi_{\mathbf{k}\uparrow\downarrow}=\sum_{l=0}^{J}\lambda_l^{\frac12}\bar\phi^l_{\uparrow\downarrow}\,\bar y_l(\mathbf{k}')\,, \tag{I.40}$$

$$\Phi_{\mathbf{k}\sigma\sigma} = \sum_{l=0}^{J} \lambda_l^{\frac{1}{2}} \phi^l_{\sigma\sigma} \left[y_l(\mathbf{k}') - y_l(-\mathbf{k}') \right], \quad \bar{\Phi}_{\mathbf{k}\sigma\sigma} = \sum_{l=0}^{J} \lambda_l^{\frac{1}{2}} \bar{\phi}^l_{\sigma\sigma} \left[\bar{y}_l(\mathbf{k}') - \bar{y}_l(-\mathbf{k}') \right] \tag{I.41}$$

and $\Omega^{\pm}_{\mathbf{k}}$ are the solutions of the quadratic equation

$$\begin{aligned} \Omega^2 - \Big(& \bar{\Phi}_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow} + \bar{\Phi}_{-\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} + \Phi_{\mathbf{k}\uparrow\uparrow}\bar{\Phi}_{\mathbf{k}\uparrow\uparrow} + \Phi_{\mathbf{k}\downarrow\downarrow}\bar{\Phi}_{\mathbf{k}\downarrow\downarrow} \Big)\Omega + \Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar{\Phi}_{\mathbf{k}\uparrow\downarrow}\bar{\Phi}_{-\mathbf{k}\uparrow\downarrow} \\ & + \bar{\Phi}_{\mathbf{k}\uparrow\uparrow}\bar{\Phi}_{\mathbf{k}\downarrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} + \Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar{\Phi}_{\mathbf{k}\uparrow\uparrow}\bar{\Phi}_{\mathbf{k}\downarrow\downarrow} + \bar{\Phi}_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\bar{\Phi}_{-\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} = 0 \end{aligned} \tag{I.42}$$

Observe that $\bar{\Phi}_{\mathbf{k}\sigma\tau}$ is not necessarily the complex conjugate of $\Phi_{\mathbf{k}\sigma\tau}$, depending on the signs of the coupling constants λ_l. The effective potential V_β is given by

$$V_\beta(\phi) = \sum_{l=0}^{J} \left(|\phi^l_{\uparrow\downarrow}|^2 + |\phi^l_{\uparrow\uparrow}|^2 + |\phi^l_{\downarrow\downarrow}|^2 \right) - \sum_{\epsilon\in\{+,-\}} \int_{M_\omega} \tfrac{d^d\mathbf{k}}{(2\pi)^d} \, \tfrac{1}{\beta} \log \left[\frac{\cosh(\frac{\beta}{2}\sqrt{e^2_{\mathbf{k}} + \Omega^\epsilon_{\mathbf{k}}})}{\cosh \frac{\beta}{2} e_{\mathbf{k}}} \right] \tag{I.43}$$

For a pure even interaction, that is, if $\lambda_\ell = 0$ for all odd angular momentum ℓ, one has, if $y_l(-\mathbf{k}) = (-1)^l y_l(\mathbf{k})$, $\Phi_{\mathbf{k}\uparrow\uparrow} = \Phi_{\mathbf{k}\downarrow\downarrow} = 0$ and the expectation value simplifies to

$$\langle \bar{\psi}_{p,\sigma}\psi_{p',\sigma} \rangle_{\beta,L} = \beta L^d \delta_{p,p'} \, \frac{\int \frac{-ip_0 - e_{\mathbf{p}}}{p_0^2 + e^2_{\mathbf{p}} + \Phi_{\mathbf{p}\uparrow\downarrow}\bar{\Phi}_{\mathbf{p}\uparrow\downarrow}} e^{-\beta L^d V_\beta(\phi_{\uparrow\downarrow})} \prod_{l=0}^{J} du^l_{\uparrow\downarrow} dv^l_{\uparrow\downarrow}}{\int e^{-\beta L^d V_\beta(\phi_{\uparrow\downarrow})} \prod_{l=0}^{J} du^l_{\uparrow\downarrow} dv^l_{\uparrow\downarrow}} \tag{I.44}$$

with an effective potential

$$V_\beta(\phi_{\uparrow\downarrow}) = \sum_{l=0}^{J} |\phi^l_{\uparrow\downarrow}|^2 - \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d} \, \tfrac{1}{\beta} \log \left[\frac{\cosh(\frac{\beta}{2}\sqrt{e^2_{\mathbf{k}} + \Phi_{\mathbf{k}\uparrow\downarrow}\bar{\Phi}_{\mathbf{k}\uparrow\downarrow}})}{\cosh \frac{\beta}{2} e_{\mathbf{k}}} \right]^2 \tag{I.45}$$

Recall that the free two point function is given by $\langle \bar{\psi}_{p,\sigma}\psi_{p',\sigma} \rangle_{\beta,L} = \beta L^d \delta_{p,p'} 1/(ip_0 - e_{\mathbf{p}}) = -\beta L^d \delta_{p,p'} (ip_0 + e_{\mathbf{p}})/(p_0^2 + e^2_{\mathbf{p}})$. In particular, for a delta function interaction one obtains the two dimensional integral representation

$$\langle \bar{\psi}_{p,\sigma}\psi_{p',\sigma} \rangle_{\beta,L} = -\beta L^d \delta_{p,p'} \, \frac{\int \frac{ip_0 + e_{\mathbf{p}}}{p_0^2 + e^2_{\mathbf{p}} + \lambda(u^2+v^2)} \, e^{-\beta L^d V_\beta(u,v)} du dv}{\int e^{-\beta L^d V_\beta(u,v)} du dv} \tag{I.46}$$

where

$$V_\beta(u,v) = u^2 + v^2 - \int_{M_\omega} \tfrac{d^d\mathbf{k}}{(2\pi)^d} \tfrac{1}{\beta} \log \left[\frac{\cosh(\frac{\beta}{2}\sqrt{e^2_{\mathbf{k}} + \lambda(u^2+v^2)})}{\cosh \frac{\beta}{2} e_{\mathbf{k}}} \right]^2 \tag{I.47}$$

A positive λ corresponds to an attractive interaction.

Since the only place where the volume L^d shows up in the integral representations (I.44,46) is the prefactor in the exponential $e^{-\beta L^d V(\phi)}$, the computation of the infinite volume limit comes down to the determination of the global minimum of the real part of the effective potential V as a function of the $\phi^l_{\sigma\tau}$'s.

figure 1

For repulsive coupling $\lambda < 0$ V_β (I.47) has a global minimum at $u = v = 0$ which results in

$$\lim_{L\to\infty} \frac{e^{-\beta L^d V_\beta(u,v)}dudv}{\int e^{-\beta L^d V_\beta(u,v)}dudv} = \delta(u)\delta(v) \tag{I.48}$$

and

$$\lim_{L\to\infty} \langle\bar\psi_{p,\sigma}\psi_{p',\sigma}\rangle_{\beta,L} = -\beta\delta_{p_0,p_0'}\delta(\mathbf{p}-\mathbf{p}')\,\tfrac{ip_0+e_{\mathbf{p}}}{p_0^2+e_{\mathbf{p}}^2} \tag{I.49}$$

For attractive coupling $\lambda > 0$ and sufficiently small $T = \frac{1}{\beta}$ the effective potential (I.47) has the form of a mexican hat. This results in

$$\lim_{L\to\infty} \langle\bar\psi_{p,\sigma}\psi_{p',\sigma}\rangle_{\beta,L} = -\beta\delta_{p_0,p_0'}\delta(\mathbf{p}-\mathbf{p}')\,\tfrac{ip_0+e_{\mathbf{p}}}{p_0^2+e_{\mathbf{p}}^2+\Delta^2} \tag{I.50}$$

where $\Delta^2 = \lambda\rho_0^2 \sim e^{-\frac{1}{\lambda}}$ and $\rho_0^2 = u_0^2 + v_0^2$ is the value where V_β takes its global minimum which lies on a circle in the u, v plane.

The expectation values $\langle\psi\psi\rangle$ and $\langle\bar\psi\bar\psi\rangle$ can also be computed. To make them nonzero, we introduce a small external field $r = |r|e^{i\alpha}$. That is, we substitute $\mathcal{V}_{\rm BCS}$ by (we do not consider here $\langle a_\uparrow a_\uparrow\rangle$ expectations)

$$\mathcal{V}_{{\rm BCS},r} = \mathcal{V}_{\rm BCS} + \tfrac{1}{\beta L^d}\sum_k [r\,\psi_{k\uparrow}\psi_{-k\downarrow} - \bar r\,\bar\psi_{k\uparrow}\bar\psi_{-k\downarrow}] \tag{I.51}$$

One obtains again a finite-dimensional integral representation. For a delta function interaction, that is for $J = 0$ in (I.37), it reduces to a two dimensional integral:

$$\langle\bar\psi_{p,\uparrow}\bar\psi_{-p',\downarrow}\rangle_{\beta,L} = \beta L^d\delta_{p,p'}\,\frac{\int \frac{-i\sqrt{\lambda}\,e^{i\alpha}(u+iv)}{p_0^2+e_{\mathbf{p}}^2+\lambda(u^2+v^2)}\,e^{-\beta L^d V_{\beta,r}(u,v)}dudv}{\int e^{-\beta L^d V_{\beta,r}(u,v)}dudv} \tag{I.52}$$

where

$$V_{\beta,r}(u,v) = u^2 + \left(v + \frac{|r|}{\sqrt{\lambda}}\right)^2 - \frac{2}{\beta}\int_M \frac{d^d\mathbf{k}}{(2\pi)^d} \log\left[\frac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\lambda(u^2+v^2)})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right] \tag{I.53}$$

Consider an attractive $\lambda > 0$ and sufficiently small $T = \frac{1}{\beta}$ such that the global minimum of the effective potential moves away from zero. For $r = 0$, the global minimum of $V_{\beta,r}$ is degenerated and lies on a circle in the u, v plane. In particular, $V_{\beta,0}$ is an even function of u and v and $\langle\bar{\psi}_{p\uparrow}\psi_{-p\downarrow}\rangle$ vanishes by symmetry.

figure 2

For $r \neq 0$, $V_{\beta,r}$ has a unique global minimum at $(u,v) = (0, v_0)$ where v_0 is given by the negative solution of

$$v_0\left\{\lambda\int_M \frac{d^d\mathbf{k}}{(2\pi)^d}\frac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\lambda v_0^2})}{2\sqrt{e_{\mathbf{k}}^2+\lambda v_0^2}} - 1\right\} = 2|r| \tag{I.54}$$

which, in the limit $r \to 0$, becomes the BCS equation for $|\Delta|^2 = \lambda v_0^2$. Thus

$$\lim_{|r|\to 0}\lim_{L\to\infty}\frac{e^{-\beta L^d V_{\beta,r}(u,v)}}{\int e^{-\beta L^d V_{\beta,r}(u,v)}dudv} = \lim_{|r|\to 0}\lim_{L\to\infty}\frac{e^{-\beta L^d[V_{\beta,r}(u,v)-V_{\beta,r}(0,v_0)]}}{\int e^{-\beta L^d[V_{\beta,r}(u,v)-V_{\beta,r}(0,v_0)]}dudv} = \delta(u)\delta\left(v + \frac{|\Delta|}{\sqrt{\lambda}}\right) \tag{I.55}$$

and $\langle\bar{\psi}_{p\uparrow}\bar{\psi}_{-p\downarrow}\rangle$ becomes nonzero.

For repulsive $\lambda < 0$, $V_{\beta,r}$ is complex and the real part of $V_{\beta,r}$ has a unique global minimum at $(u,v) = (0,0)$ which results in $\lim_{|r|\to 0}\lim_{L\to\infty}\frac{e^{-\beta L^d V_{\beta,r}(u,v)}}{\int e^{-\beta L^d V_{\beta,r}(u,v)}dudv} = \delta(u)\delta(v)$ and $\lim_{|r|\to 0}\lim_{L\to\infty}\langle\bar{\psi}_{p\uparrow}\bar{\psi}_{-p\downarrow}\rangle = 0$.

These results of course are also obtained if one applies the usual mean field formalism which is based on approximating the quartic Hamiltonian by a quadratic expression. However, the situation is different if the electron-electron interaction contains higher angular momentum terms and the space dimension is 3. In that case the standard mean field formalism [AB,BW] predicts an angle dependent gap but, if $e_{\mathbf{k}}$ has $SO(3)$ symmetry, this is not the case.

Suppose that $\lambda_\ell > 0$ is attractive and

$$U(\mathbf{k}-\mathbf{p}) = \lambda_\ell \sum_{m=-\ell}^{\ell} \bar{Y}_{\ell m}(\mathbf{k}') Y_{\ell m}(\mathbf{p}') \tag{I.56}$$

Then the Anderson Balian Werthammer mean field formalism gives

$$\lim_{L\to\infty} \tfrac{1}{L^d}\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}\rangle = \tfrac{1}{2}\left(1 - e_{\mathbf{k}}\left[\tfrac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}})}{\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}}}\right]_{\sigma\sigma}\right) \tag{I.57}$$

where the 2×2 matrix $\Delta_{\mathbf{k}}$, $\Delta^T_{\mathbf{k}} = -\Delta_{-\mathbf{k}}$, is a solution of the gap equation

$$\Delta_{\mathbf{p}} = \int_{M_\omega} \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, U(\mathbf{p}'-\mathbf{k}')\, \Delta_{\mathbf{k}} \tfrac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}})}{2\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}}} \tag{I.58}$$

In 3 dimensions, it has been proven [FKT2] that for all $\ell \geq 2$ (IV.58) does not have unitary isotropic ($\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}} = const\, Id$) solutions. That is, the gap in (I.57) is angle dependent. However in Theorem III.3 it is shown that, for even ℓ,

$$\lim_{L\to\infty} \tfrac{1}{\beta L^d}\langle \bar{\psi}_{k\sigma}\psi_{k\sigma}\rangle_{\beta,L} = \int_{S^2} \tfrac{ik_0+e_{\mathbf{k}}}{k_0^2+e_{\mathbf{k}}^2+\lambda_\ell\rho_0^2|\sum_m \alpha^0_m Y_{\ell m}(\mathbf{x})|^2}\,\tfrac{d\Omega(\mathbf{x})}{4\pi} \tag{I.59}$$

which gives, using (I.25),

$$\lim_{L\to\infty} \tfrac{1}{L^d}\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}\rangle = \int_{S^2} \tfrac{1}{2}\left(1 - e_{\mathbf{k}}\tfrac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+|\Delta(\mathbf{x})|^2})}{\sqrt{e_{\mathbf{k}}^2+|\Delta(\mathbf{x})|^2}}\right)\tfrac{d\Omega(\mathbf{x})}{4\pi} \tag{I.60}$$

if $\Delta(\mathbf{x}) = \lambda_\ell^{\frac{1}{2}}\rho_0 \sum_m \alpha^0_m Y_{\ell m}(\mathbf{x})$ and ρ_0 and α^0 are values at the global minimum. The point is that for $SO(3)$ symmetric $e_{\mathbf{k}}$ also the effective potential has $SO(3)$ symmetry which means that also the global minimum has $SO(3)$ symmetry. Since in the infinite volume limit the integration variables are forced to be at the global minimum, the integral over the sphere in (I.59,60) is the averaging over all global minima.

There is some physics literature [B,BZT,BR,G,Ha,W] which investigates the relation between the reduced but still quartic and not solvable BCS Hamiltonian

$$H_{\mathrm{BCS}} = H_0 + \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}}\sum_{\mathbf{k},\mathbf{p}} U(\mathbf{k}-\mathbf{p})\, a^+_{\mathbf{k}\sigma}a^+_{-\mathbf{k}\tau}a_{\mathbf{p}\sigma}a_{-\mathbf{p}\tau} \tag{I.61}$$

and the quadratic, explicitly diagonalizable mean field Hamiltonian

$$\begin{aligned} H_{\mathrm{MF}} = H_0 + \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}}\sum_{\mathbf{k},\mathbf{p}} U(\mathbf{k}-\mathbf{p}) \Big(& a^+_{\mathbf{k}\sigma}a^+_{-\mathbf{k}\tau}\langle a_{\mathbf{p}\sigma}a_{-\mathbf{p}\tau}\rangle + \langle a^+_{\mathbf{k}\sigma}a^+_{-\mathbf{k}\tau}\rangle a_{\mathbf{p}\sigma}a_{-\mathbf{p}\tau} \\ & - \langle a^+_{\mathbf{k}\sigma}a^+_{-\mathbf{k}\tau}\rangle\langle a_{\mathbf{p}\sigma}a_{-\mathbf{p}\tau}\rangle\Big) \end{aligned} \tag{I.62a}$$

where the numbers $\langle a_{\mathbf{p}\sigma} a_{-\mathbf{p}\tau}\rangle$ have to be determined by the condition

$$\langle a_{\mathbf{p}\sigma} a_{-\mathbf{p}\tau}\rangle = \frac{Tr\; e^{-\beta H_{\rm MF}} a_{\mathbf{p}\sigma} a_{-\mathbf{p}\tau}}{Tr\; e^{-\beta H_{\rm MF}}} \tag{I.62b}$$

which is equivalent to the gap equation (I.58) if one defines $\Delta_{\mathbf{p}\sigma\tau} = \frac{1}{L^d}\sum_{\mathbf{k}} U(\mathbf{p}-\mathbf{k}) \frac{1}{L^d}\langle a_{\mathbf{k}\sigma} a_{-\mathbf{k}\tau}\rangle$. One has, if $H' := H_{\rm BCS} - H_{\rm MF}$,

$$H' = \tfrac{1}{L^{3d}} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{\mathbf{k},\mathbf{p}} U(\mathbf{k}-\mathbf{p}) \Big(a^+_{\mathbf{k}\sigma} a^+_{-\mathbf{k}\tau} - \langle a^+_{\mathbf{k}\sigma} a^+_{-\mathbf{k}\tau}\rangle \Big) \Big(a_{\mathbf{p}\sigma} a_{-\mathbf{p}\tau} - \langle a_{\mathbf{p}\sigma} a_{-\mathbf{p}\tau}\rangle \Big) \tag{I.63}$$

It is claimed that, in the infinite volume limit, the correlation functions of both models should coincide. More precisely, it is claimed that

$$\lim_{L\to\infty} \tfrac{1}{L^d} \log \tfrac{Tr\, e^{-\beta H_{\rm BCS}}}{Tr\, e^{-\beta H_{\rm MF}}} \tag{I.64}$$

vanishes. To this end it is argued that each order of perturbation theory (with respect to H') of $Tr\, e^{-\beta(H_{\rm MF}+H')}/Tr\, e^{-\beta H_{\rm MF}}$ is finite as the volume goes to infinity. The Haag paper argues that spatial averages of field operators like $\frac{1}{L^d}\int_{[0,L]^d} d^d\mathbf{x} a_\uparrow(\mathbf{x}) a_\downarrow(\mathbf{x})$ may be substituted by numbers in the infinite volume limit, since commutators with them have an extra one over volume factor, but there is no rigorous controll of the error. At least for a more complicated electron-electron interaction (I.56), this reasoning cannot be correct in view of (I.57-60).

Namely, consider the $\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}\rangle_{\beta,L}$ expectation. In terms of Grassmann variables it is given by (I.25,26). Assume first a delta function interaction $U(\mathbf{k}-\mathbf{p}) = \lambda$. For the full model (I.3), by making a Hubbard Stratonovich transformation, (I.26) can be rewritten as

$$\langle \bar\psi_{p\sigma} \psi_{p\sigma}\rangle = \frac{\int F(\phi)\, e^{-V(\phi)} \prod_{q_0\mathbf{q}} d\phi_{q_0\mathbf{q}} d\bar\phi_{q_0\mathbf{q}}}{\int e^{-V(\phi)} \prod_{q_0\mathbf{q}} d\phi_{q_0\mathbf{q}} d\bar\phi_{q_0\mathbf{q}}} \tag{I.65}$$

where the integrand is given by

$$F(\phi) = \frac{\frac{\partial}{\partial s_{p\sigma}}_{|s=0} \det \begin{bmatrix} A + S_\uparrow & i(\frac{\lambda}{\beta L^d})^{\frac12}\,\phi \\ i(\frac{\lambda}{\beta L^d})^{\frac12}\,\bar\phi & \bar A + S_\downarrow \end{bmatrix}}{\det \begin{bmatrix} A & i(\frac{\lambda}{\beta L^d})^{\frac12}\,\phi \\ i(\frac{\lambda}{\beta L^d})^{\frac12}\,\bar\phi & \bar A \end{bmatrix}} \tag{I.66}$$

and the effective potential reads

$$V(\phi) = \sum_{q_0\mathbf{q}} |\phi_{q_0\mathbf{q}}|^2 - \log \frac{\det \begin{bmatrix} A & i(\frac{\lambda}{\beta L^d})^{\frac12}\,\phi \\ i(\frac{\lambda}{\beta L^d})^{\frac12}\,\bar\phi & \bar A \end{bmatrix}}{\det \begin{bmatrix} A & 0 \\ 0 & \bar A \end{bmatrix}} \tag{I.67}$$

Here A and S_σ are diagonal matrices with entries $A_k = ik_0 - e_{\mathbf{k}}$ and $s_{k\sigma}$ respectively and ϕ is a short notation for the matrix $(\phi_{k-p})_{k,p}$ where $k = (k_0, \mathbf{k}) \in \frac{\pi}{\beta}(2\mathbf{Z}+1) \times M_\omega$. Of course, for example $\det \left[\begin{smallmatrix} A & 0 \\ 0 & \bar{A} \end{smallmatrix}\right] = \Pi_k (k_0^2 + e_{\mathbf{k}}^2)$ does not make sense, but the quotients in (I.66,67) are well defined.

The correlation functions of the reduced, but still quartic BCS Hamiltonian (I.61) are obtained from (I.65-67) by first assuming that the global minimum $\phi_q^{\min}$ of $V(\phi)$ is proportional to $\delta_{\mathbf{q},\mathbf{0}}\phi_{q_0}$ and second by suppressing the quantum fluctuations around $\phi_{\mathbf{q}q_0}^{\min} = 0$ for all $\mathbf{q} \neq 0$. That is, for the model (I.61) one finds

$$\langle \bar{\psi}_{p\sigma} \psi_{p\sigma} \rangle = \frac{\int F_{\mathbf{p}}(\phi)\, e^{-L^d V(\phi)} \prod\limits_{q_0} d\phi_{q_0} d\bar{\phi}_{q_0}}{\int e^{-L^d V(\phi)} \prod\limits_{q_0} d\phi_{q_0} d\bar{\phi}_{q_0}} \tag{I.68}$$

where the integrand is given by

$$F_{\mathbf{p}}(\phi) = \frac{\frac{\partial}{\partial s_{p_0\sigma}}_{|s=0} \det \begin{bmatrix} A_{\mathbf{p}} + S_{\mathbf{p}\uparrow} & i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \phi \\ i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \bar{\phi} & \bar{A}_{\mathbf{p}} + S_{\mathbf{p}\downarrow} \end{bmatrix}}{\det \begin{bmatrix} A_{\mathbf{p}} & i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \phi \\ i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \bar{\phi} & \bar{A}_{\mathbf{p}} \end{bmatrix}} \tag{I.69}$$

and the effective potential reads

$$V(\phi) = \sum_{q_0} |\phi_{q_0}|^2 - \tfrac{1}{L^d} \sum_{\mathbf{k}} \log \frac{\det \begin{bmatrix} A_{\mathbf{k}} & i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \phi \\ i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \bar{\phi} & \bar{A}_{\mathbf{k}} \end{bmatrix}}{\det \begin{bmatrix} A_{\mathbf{k}} & 0 \\ 0 & \bar{A}_{\mathbf{k}} \end{bmatrix}} \tag{I.70}$$

The volume factor $L^{-\frac{d}{2}}$ in the determinant in (I.66,67) has been transformed away by a substitution of variables such that it shows up in the exponent in front of the effective potential in (I.68). The matrices in (I.69,70) are labelled only by the k_0, p_0 variables, that is, $A_{\mathbf{k}}$ is the diagonal matrix with entries $A_{\mathbf{k},k_0} = ik_0 - e_{\mathbf{k}}$ and ϕ is a short notation for the matrix $(\phi_{k_0-p_0})_{k_0,p_0}$.

Contrary to the full model, the volume dependence of the model (I.61) or (I.68-70) is such that in the infinite volume limit the integration variables ϕ_{q_0} in (I.68) are forced to be at the global minimum of (I.70).

The model discussed in this paper (that is, only the BCS part) is obtained from (I.68-70) by assuming that the global minimum of (I.70) is proportional to $\delta_{q_0,0}$. In that case, the only integration variable which is left in (I.68) is the $q_0 = 0$ mode $\phi = \phi_0$ and the expressions (I.68-70) reduce to the integral representation (I.46,47).

Now assume that the elctron electron interaction is given by (I.56) and suppose for simplicity that ℓ is even which suppresses $\uparrow\uparrow\uparrow\uparrow$ and $\downarrow\downarrow\downarrow\downarrow$ contributions in (I.3) and (I.61). In that case the model (I.61) gives

$$\langle\bar{\psi}_{p\sigma}\psi_{p\sigma}\rangle = \frac{\int F_{\mathbf{p}}(\phi)\, e^{-L^d V(\phi)} \prod\limits_{m=-\ell}^{\ell} \prod\limits_{q_0} d\phi^m_{q_0} d\bar{\phi}^m_{q_0}}{\int e^{-L^d V(\phi)} \prod\limits_{m=-\ell}^{\ell} \prod\limits_{q_0} d\phi^m_{q_0} d\bar{\phi}^m_{q_0}} \tag{I.71}$$

where the integrand is given by

$$F_{\mathbf{p}}(\phi) = \frac{\frac{\partial}{\partial s_{p_0\sigma}}_{|s=0} \det \begin{bmatrix} A_{\mathbf{p}} + S_{\mathbf{p}\uparrow} & i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \Phi_{\mathbf{p}} \\ i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \bar{\Phi}_{\mathbf{p}} & \bar{A}_{\mathbf{p}} + S_{\mathbf{p}\downarrow} \end{bmatrix}}{\det \begin{bmatrix} A_{\mathbf{p}} & i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \Phi_{\mathbf{p}} \\ i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \bar{\Phi}_{\mathbf{p}} & \bar{A}_{\mathbf{p}} \end{bmatrix}} \tag{I.72}$$

and the effective potential reads

$$V(\phi) = \sum_{m=-\ell}^{\ell} \sum_{q_0} |\phi^m_{q_0}|^2 - \tfrac{1}{L^d} \sum_{\mathbf{k}} \log \frac{\det \begin{bmatrix} A_{\mathbf{k}} & i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \Phi_{\mathbf{k}} \\ i(\frac{\lambda}{\beta})^{\frac{1}{2}}\, \bar{\Phi}_{\mathbf{k}} & \bar{A}_{\mathbf{k}} \end{bmatrix}}{\det \begin{bmatrix} A_{\mathbf{k}} & 0 \\ 0 & \bar{A}_{\mathbf{k}} \end{bmatrix}} \tag{I.73}$$

Here $\Phi_{\mathbf{k}}$ denotes the matrix with entries $\Phi_{\mathbf{k},p_0p_0'} = \sum_{m=-\ell}^{\ell} \phi^m_{p_0-p_0'} Y_{\ell m}(\mathbf{k})$, labelled by the p_0, p_0' variables. The model discussed in this paper is obtained from (I.71-73) by assuming that the global minimum of (I.73) has only nonzero $\phi^m_{q_0=0}$ modes. In that case (I.71-73) reduce to the integral representation (I.44,45) which further can be reduced, by Theorem III.3, to (I.59). However, the argument used in Theorem III.3 that the $\langle\bar{\psi}_{p\sigma}\psi_{p\sigma}\rangle$ expectation has to be $SO(3)$ invariant if $e_{\mathbf{k}}$ is $SO(3)$ invariant still applies to (I.71), since (I.73) is invariant under simultanious transformation of the $\phi^m_{q_0}$'s to $\sum_{m'} U(R)_{mm'}\phi^{m'}_{q_0}$ where $U(R)$ is the unitary representation of $SO(3)$ given by $Y_{\ell m}(R\mathbf{k}) = \sum_{m'} U(R)_{mm'} Y_{\ell m'}(\mathbf{k})$.

That is, if the BCS equation (I.58) or (I.62b) of the quadratic mean field model (I.62) has a solution such that $|\Delta_{\mathbf{k}}|^2$ is not $SO(3)$ invariant (and, by [FKT2], this is necessarily the case for any nonzero solution for $d = 3$ and $\ell \geq 2$), then the $\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}\rangle$ expectation of the quadratic mean field model (I.62) does not coincide with the corresponding expectation of the quartic reduced BCS Hamiltonian (I.61).

In [AB], Anderson and Brinkmann used the quadratic mean field Hamiltonian with an $\ell = 1$ interaction to describe the properties of superfluid Helium 3. The basic quantities in their analysis are the $\langle a_{\mathbf{k}\sigma} a_{-\mathbf{k}\tau}\rangle$ expectations or the matrix $\Delta_{\mathbf{k}\sigma\tau}$ which is obtained as a solution of the gap equation of the quadratic model. In view of the discussion above, one may regard as the more natural approach to take the quartic BCS Hamiltonian (I.61), to add in a symmetry breaking term which breaks the U(1) particle symmetry as well as the spatial SO(3) symmetry and then to compute (in the approximation $\phi^m_{q_0} = \delta_{q_0,0}\phi^m$) the infinite volume limit followed by the limit symmetry breaking term → 0. In particular, besides the usual U(1) symmetry breaking one may expect SO(3) symmetry breaking in the sense that probably also for the $\langle a^+ a\rangle$ expectations the above two limits do not commute. That is, if B denotes the SO(3) symmetry breaking term, whereas $\lim_{L\to\infty}\lim_{B\to 0}\langle a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}\rangle$ has SO(3) symmetry, $\lim_{B\to 0}\lim_{L\to\infty}\langle a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}\rangle$ may have not. However, to what extend the quantities of the [AB] paper are related to these expectations is not clear. For example, for $\ell \geq 2$ ([AB] has $\ell = 1$, but still uses the quadratic mean field formalism) any nontrivial solution of the quadratic model (I.62) is necessarily anisotropic, without any external SO(3) symmetry breaking field at all. The quantities $\lim_{B\to 0}\lim_{L\to\infty}\langle a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}\rangle$ for the quartic model (I.61) or the model discussed in this paper of course would depend on the direction of B, if they become anisotropic. A more careful analysis of this question we defer to another paper.

Acknowledgements

I am grateful to Horst Knörrer and Eugene Trubowitz and to the Forschungsinstitut für Mathematik at ETH Zürich for the hospitality and the support during the summer 1996. Furthermore I would like to thank Joel Feldman who made it possible for me to visit the University of British Columbia in Vancouver in the academic year 1996/97.

II. Solution of the Model in the Forward, Exchange and BCS Approximation

Let $H = H_0 + H_{\rm int}$ where

$$H_0 = \tfrac{1}{L^d}\sum_{\sigma\in\{\uparrow,\downarrow\}}\sum_{\mathbf{k}\in M} e_{\mathbf{k}}\, a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma} \tag{I.2}$$

and

$$H_{\rm int} = \tfrac{1}{L^{3d}}\sum_{\sigma,\tau\in\{\uparrow,\downarrow\}}\sum_{\mathbf{k},\mathbf{p},\mathbf{q}} U(\mathbf{k}-\mathbf{p})\, a^+_{\mathbf{k},\sigma}a_{\mathbf{p},\sigma}a^+_{\mathbf{q}-\mathbf{k},\tau}a_{\mathbf{q}-\mathbf{p},\tau} \tag{I.3}$$

Then

$$Z = \frac{Tr\, e^{-\beta(H_0+H_{int})}}{Tr\, e^{-\beta H_0}} = \int e^{-\mathcal{V}_{\rm int}(\psi,\bar\psi)}d\mu_C(\psi,\bar\psi) \tag{I.23}$$

where the exponent in the fermionic integral is given by ($\kappa = \beta L^d$)

$$\mathcal{V}_{\rm int}(\psi) = \tfrac{1}{\kappa^3} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{k,p,q} U(\mathbf{k}-\mathbf{p})\, \bar\psi_{k,\sigma}\psi_{p,\sigma}\bar\psi_{q-k,\tau}\psi_{q-p,\tau} \tag{I.24}$$

and $k = (k_0, \mathbf{k}) \in \frac{\pi}{\beta}(2\mathbf{Z}+1) \times M$.

The forward, exchange and BCS approximation is obtained by restricting the above sum to the following terms

$$\text{forward}: \quad \delta_{k,p}\ , \qquad\quad \text{exchange}: \quad \delta_{k,q-p}\ , \qquad\quad \text{BCS}: \quad \delta_{q,0}\ . \tag{I.35}$$

We consider the approximation $\mathcal{V}_{\rm int} \approx \mathcal{V}_{\rm forw} + \mathcal{V}_{\rm ex} + \mathcal{V}_{\rm BCS} \equiv \mathcal{U}(\psi, \bar\psi)$. That is,

$$\mathcal{U}(\psi,\bar\psi) = \tfrac{1}{\kappa^3} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{k,p,q} U(\mathbf{k}-\mathbf{p})\big[\delta_{k,p} + \delta_{k,q-p} + \delta_{q,0}\big]\, \bar\psi_{k,\sigma}\psi_{p,\sigma}\bar\psi_{q-k,\tau}\psi_{q-p,\tau} \tag{II.1}$$

and

$$U(\mathbf{k}-\mathbf{p}) = -\begin{cases} \dfrac{1}{2}\displaystyle\sum_{\ell=-j}^{j} \lambda_{|\ell|} e^{i\ell\varphi_{\mathbf{k}}} e^{-i\ell\varphi_{\mathbf{p}}} + \dfrac{\lambda_0}{2} & \text{if } d=2 \\ \displaystyle\sum_{\ell=0}^{j}\sum_{m=-\ell}^{\ell} \lambda_\ell \bar Y_{\ell m}\left(\mathbf{k}'\right) Y_{\ell m}\left(\mathbf{p}'\right) & \text{if } d=3 \end{cases} =: \ -\sum_{l=0}^{J} \lambda_l\, y_l(\mathbf{k}')\, \bar y_l(\mathbf{p}') \tag{I.37}$$

and we abbreviated

$$\kappa = \beta L^d \tag{II.2}$$

This model can be solved explicitly. Before we write down the general solution we shortly indicate the computation for $\mathcal{V}_{\rm int} \approx \mathcal{V}_{\rm BCS}$ and $U(\mathbf{k}-\mathbf{p}) = -\lambda$. In that case it comes down to the usual effective potential computation with ‘constant ϕ’. That is, using the identity ($\phi = u + iv$, $\bar\phi = u - iv \in \mathbf{C}$)

$$e^{2ab} = \tfrac{1}{2\pi} \int_{\mathbf{R}^2} e^{a\phi + b\bar\phi} e^{-\frac{1}{2}|\phi|^2} dudv \tag{II.3}$$

one obtains

$$\begin{aligned} Z &= \int e^{-\left(\frac{\lambda}{\kappa}\right)^{\frac12} \frac{1}{\kappa}\sum_k \bar\psi_{k\uparrow}\bar\psi_{-k\downarrow} \times \left(\frac{\lambda}{\kappa}\right)^{\frac12}\frac{1}{\kappa}\sum_p \psi_{p\uparrow}\psi_{-p\downarrow}} d\mu_C(\psi,\bar\psi) \\ &= \int_{\mathbf{R}^2}\int e^{i\left(\frac{\lambda}{\kappa}\right)^{\frac12}\phi\,\frac{1}{\kappa}\sum_k \psi_{k\uparrow}\psi_{-k\downarrow} + i\left(\frac{\lambda}{\kappa}\right)^{\frac12}\bar\phi\,\frac{1}{\kappa}\sum_k \bar\psi_{k\uparrow}\bar\psi_{-k\downarrow}} d\mu_C(\psi,\bar\psi) \tfrac{1}{2\pi} e^{-\frac12|\phi|^2} dudv \\ &= \int_{\mathbf{R}^2} \prod_k \tfrac{1}{k_0^2+e_{\mathbf{k}}^2} \det\begin{pmatrix} ik_0 - e_{\mathbf{k}} & i\lambda^{\frac12}\bar\phi \\ i\lambda^{\frac12}\phi & -ik_0 - e_{\mathbf{k}} \end{pmatrix} \tfrac{\kappa}{\pi} e^{-\kappa|\phi|^2} dudv \\ &= \tfrac{\kappa}{\pi}\int_{\mathbf{R}^2} e^{-\kappa V_\beta(u,v)} dudv \end{aligned} \tag{II.4}$$

In the following theorem the ξ variables take care of the exchange contributions, the w variable makes the forward contribution quadratic in the fermion fields and the ϕ variables, as in (II.3,4) above, sum up BCS contributions.

Theorem II.1: *Let $\mathcal{U}(\psi,\bar\psi)$ be given by (II.1) and let*

$$Z(\beta,L,\{s_{k,\sigma}\}) = \int e^{-\mathcal{U}(\psi,\bar\psi)+\frac{1}{\kappa}\sum_{k,\sigma} s_{k,\sigma}\bar\psi_{k,\sigma}\psi_{k,\sigma}}\, d\mu_C(\psi,\bar\psi) \tag{II.5}$$

$$\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle = \kappa\,\frac{\partial}{\partial s_{p\sigma}}_{|s=0}\log Z(\beta,L,\{s_{k\sigma}\}) \tag{II.6}$$

Let $w\in\mathbf{R}$, $\xi^l_{\sigma\tau} = a^l_{\sigma\tau}+ib^l_{\sigma\tau}$, $\phi^l_{\sigma\tau} = u^l_{\sigma\tau}+iv^l_{\sigma\tau}\in\mathbf{C}$ for $0\le l\le J$, $(\sigma\tau)\in\{\uparrow\uparrow,\downarrow\downarrow,\uparrow\downarrow\}$ and define the fields

$$\Xi_{\mathbf{k}\uparrow\downarrow} = \sum_{l=0}^{J}(2\lambda_l)^{\frac12}\xi^l_{\uparrow\downarrow}\,y_l(\mathbf{k}')\,,\quad \bar\Xi_{\mathbf{k}\uparrow\downarrow} = \sum_{l=0}^{J}(2\lambda_l)^{\frac12}\bar\xi^l_{\uparrow\downarrow}\,\bar y_l(\mathbf{k}')$$

$$\Xi_{\mathbf{k}\sigma\sigma} = \sum_{l=0}^{J}\lambda_l^{\frac12}\,\xi^l_{\sigma\sigma}\,y_l(\mathbf{k}')\,,\quad \bar\Xi_{\mathbf{k}\sigma\sigma} = \sum_{l=0}^{J}\lambda_l^{\frac12}\,\bar\xi^l_{\sigma\sigma}\,\bar y_l(\mathbf{k}')\,, \tag{II.7}$$

$$\Gamma_{\mathbf{k}\uparrow} = U_0^{\frac12}w + i\Xi_{\mathbf{k}\uparrow\uparrow} + i\bar\Xi_{\mathbf{k}\uparrow\uparrow}\,,\quad \Gamma_{\mathbf{k}\downarrow} = U_0^{\frac12}w + i\Xi_{\mathbf{k}\downarrow\downarrow} + i\bar\Xi_{\mathbf{k}\downarrow\downarrow}\,, \tag{II.8}$$

$$\Phi_{\mathbf{k}\uparrow\downarrow} = \sum_{l=0}^{J}(2\lambda_l)^{\frac12}\phi^l_{\uparrow\downarrow}\,y_l(\mathbf{k}')\,,\quad \bar\Phi_{\mathbf{k}\uparrow\downarrow} = \sum_{l=0}^{J}(2\lambda_l)^{\frac12}\bar\phi^l_{\uparrow\downarrow}\,\bar y_l(\mathbf{k}') \tag{II.9}$$

$$\Phi_{\mathbf{k}\sigma\sigma} = \sum_{l=0}^{J}\lambda_l^{\frac12}\,\phi^l_{\sigma\sigma}\,[y_l(\mathbf{k}')-y_l(-\mathbf{k}')]\,,\quad \bar\Phi_{\mathbf{k}\sigma\sigma} = \sum_{l=0}^{J}\lambda_l^{\frac12}\,\bar\phi^l_{\sigma\sigma}\,[\bar y_l(\mathbf{k}')-\bar y_l(-\mathbf{k}')]\,. \tag{II.10}$$

where $U_0 = U(\mathbf{k}=\mathbf{0}) = \int d^d\mathbf{x}\,U(\mathbf{x})$. For each $k=(k_0,\mathbf{k})$, let S_k be the 8×8 skew symmetric matrix

$$S_k = -i\begin{pmatrix}
0 & -\frac1i A_{k\uparrow} & \bar\Phi_{\mathbf{k}\uparrow\downarrow} & 0 & \bar\Xi_{\mathbf{k}\uparrow\downarrow} & 0 & 0 & \bar\Phi_{\mathbf{k}\uparrow\uparrow}\\
\frac1i A_{k\uparrow} & 0 & 0 & \Phi_{\mathbf{k}\uparrow\downarrow} & 0 & -\Xi_{\mathbf{k}\uparrow\downarrow} & \Phi_{\mathbf{k}\uparrow\uparrow} & 0\\
-\bar\Phi_{\mathbf{k}\uparrow\downarrow} & 0 & 0 & -\frac1i A_{-k\downarrow} & 0 & -\bar\Phi_{\mathbf{k}\downarrow\downarrow} & \Xi_{-\mathbf{k}\uparrow\downarrow} & 0\\
0 & -\Phi_{\mathbf{k}\uparrow\downarrow} & \frac1i A_{-k\downarrow} & 0 & -\Phi_{\mathbf{k}\downarrow\downarrow} & 0 & 0 & -\bar\Xi_{-\mathbf{k}\uparrow\downarrow}\\
-\bar\Xi_{\mathbf{k}\uparrow\downarrow} & 0 & 0 & \Phi_{\mathbf{k}\downarrow\downarrow} & 0 & \frac1i A_{k\downarrow} & -\Phi_{-\mathbf{k}\uparrow\downarrow} & 0\\
0 & \Xi_{\mathbf{k}\uparrow\downarrow} & \bar\Phi_{\mathbf{k}\downarrow\downarrow} & 0 & -\frac1i A_{k\downarrow} & 0 & 0 & -\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\\
0 & -\Phi_{\mathbf{k}\uparrow\uparrow} & -\Xi_{-\mathbf{k}\uparrow\downarrow} & 0 & \Phi_{-\mathbf{k}\uparrow\downarrow} & 0 & 0 & \frac1i A_{-k\uparrow}\\
-\bar\Phi_{\mathbf{k}\uparrow\uparrow} & 0 & 0 & \bar\Xi_{-\mathbf{k}\uparrow\downarrow} & 0 & \bar\Phi_{-\mathbf{k}\uparrow\downarrow} & -\frac1i A_{-k\uparrow} & 0
\end{pmatrix}$$

where $A_{k\sigma} = a_k - s_{k\sigma} - \Gamma_{k\sigma} = ik_0 - e_{\mathbf{k}} - s_{k\sigma} - \Gamma_{k\sigma}$ and let $\mathrm{Pf}S_k$ be the Pfaffian of S_k given by Lemma II.2 below. Then:

a)

$$Z(\beta,L,\{s_{k,\sigma}\}) = \int \prod_{\substack{k_0>0\\ \mathbf{k}\in M}}\Big\{\Big(\tfrac{1}{a_k a_{-k}}\Big)^2\mathrm{Pf}S_k\Big\}\,d\nu_\kappa(w,\xi,\phi) \tag{II.11}$$

where

$$d\nu_\kappa(w,\xi,\phi) = \left(\tfrac{\kappa}{4\pi}\right)^{\frac{1}{2}} e^{-\frac{\kappa}{4}w^2} dw \prod_{l=0}^{J}\prod_{\sigma\tau}\Big\{\left(\tfrac{\kappa}{\pi}\right)^2\; e^{-\kappa(|\phi^l_{\sigma\tau}|^2+|\xi^l_{\sigma\tau}|^2)} du^l_{\sigma\tau} dv^l_{\sigma\tau} da^l_{\sigma\tau} db^l_{\sigma\tau}\Big\} \tag{II.12}$$

b)

$$\langle\bar\psi_{p\sigma}\psi_{p'\sigma}\rangle = \beta L^d \delta_{p,p'}\; \frac{\int \frac{\frac{\partial}{\partial s_{p\sigma}}\mathrm{Pf}S_p}{\mathrm{Pf}S_p} e^{-\kappa V(w,\xi,\phi)} dw \prod d\xi d\phi}{\int\;\; e^{-\kappa V(w,\xi,\phi)} dw \prod d\xi d\phi} \tag{II.13}$$

where the effective potential V *is given by*

$$V(w,\xi,\phi) = \tfrac{1}{4}w^2 + \sum_{\sigma\tau}\sum_{l=0}^{J}(|\phi^l_{\sigma\tau}|^2 + |\xi^l_{\sigma\tau}|^2) - \tfrac{1}{\beta L^d}\sum_{\mathbf{k}\in M}\log\prod_{k_0>0}\tfrac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2} \tag{II.14}$$

and $\prod d\xi d\phi = \prod_{\sigma\tau}\prod_{l=0}^{J} d\xi^l_{\sigma\tau} d\phi^l_{\sigma\tau}$*. Here* $\sigma\tau \in \{\uparrow\uparrow, \downarrow\downarrow, \uparrow\downarrow\}$*.*

Remark: The product $\prod_{k_0>0}\frac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2}$ in the effective potential (II.14) where $\mathrm{Pf}S_k = \mathrm{Pf}S_k(a_k, a_{-k})$ has to be computed according to the rule

$$\prod_{k_0>0}\frac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2} = \lim_{\substack{\epsilon\to 0\\ \epsilon<0}}\prod_{k_0>0}\frac{\mathrm{Pf}S_k(e^{ik_0\epsilon}a_k, e^{-ik_0\epsilon}a_{-k})}{a_k^2 a_{-k}^2} \tag{II.15}$$

That there are cases where it is necessary to make the phase factors explicit can be seen from the discussion given in the proof of Lemma A1 in the appendix.

Proof: Since we assume $U(\mathbf{p}-\mathbf{k}) = U(\mathbf{k}-\mathbf{p})$, one has

$$\begin{aligned}
\mathcal{U}(\psi,\bar\psi) &= \tfrac{1}{\kappa^3}\sum_{\sigma,\tau}\sum_{k,p} U(\mathbf{0})\,\bar\psi_{k,\sigma}\bar\psi_{p,\tau}\psi_{k,\sigma}\psi_{p,\tau} + \tfrac{1}{\kappa^3}\sum_{\sigma,\tau}\sum_{k,p} U(\mathbf{k}-\mathbf{p})\,\bar\psi_{p,\sigma}\bar\psi_{k,\tau}\psi_{k,\sigma}\psi_{p,\tau} \\
&\qquad + \tfrac{1}{\kappa^3}\sum_{\sigma,\tau}\sum_{k,p} U(\mathbf{k}-\mathbf{p})\,\bar\psi_{p,\sigma}\bar\psi_{-p,\tau}\psi_{k,\sigma}\psi_{-k,\tau} \\
&= -\tfrac{1}{\kappa^2}\tfrac{U_0}{\kappa}\sum_{k,p}\Big(\bar\psi_{k\uparrow}\psi_{k\uparrow} + \bar\psi_{k\downarrow}\psi_{k\downarrow}\Big)\Big(\bar\psi_{p\uparrow}\psi_{p\uparrow} + \bar\psi_{p\downarrow}\psi_{p\downarrow}\Big) \\
&\qquad + \tfrac{1}{\kappa^3}\sum_{k,p} U(\mathbf{k}-\mathbf{p})\,\Big\{\bar\psi_{p\uparrow}\psi_{p\uparrow}\bar\psi_{k\uparrow}\psi_{k\uparrow} + \bar\psi_{p\downarrow}\psi_{p\downarrow}\bar\psi_{k\downarrow}\psi_{k\downarrow} + 2\bar\psi_{p\uparrow}\psi_{p\downarrow}\bar\psi_{k\downarrow}\psi_{k\uparrow}\Big\} \\
&\qquad + \tfrac{1}{\kappa^3}\sum_{k,p} U(\mathbf{k}-\mathbf{p})\Big\{\bar\psi_{p\uparrow}\bar\psi_{-p\uparrow}\psi_{k\uparrow}\psi_{-k\uparrow} + \bar\psi_{p\downarrow}\bar\psi_{-p\downarrow}\psi_{k\downarrow}\psi_{-k\downarrow} + 2\bar\psi_{p\uparrow}\bar\psi_{-p\downarrow}\psi_{k\uparrow}\psi_{-k\downarrow}\Big\}
\end{aligned}$$

We substitute $U(\mathbf{k}-\mathbf{p}) = \sum_{l=0}^{J} \lambda_l y_l(\mathbf{k}')\bar{y}_l(\mathbf{p}')$ and use the identities

$$e^{\frac{1}{2}X^2} = \tfrac{1}{\sqrt{2\pi}} \int_{\mathbf{R}} e^{Xw} e^{-\frac{1}{2}w^2} dw$$

$$e^{-2XY} = \tfrac{1}{2\pi} \int_{\mathbf{R}^2} e^{iX\phi + iY\bar{\phi}} e^{-\frac{1}{2}|\phi|^2} du dv$$

to obtain

$$\begin{aligned}
&e^{-\mathcal{U}(\psi,\bar{\psi})} = \\
&\quad \int \exp\Big\{ \big(\tfrac{2U_0}{\kappa}\big)^{\frac{1}{2}}\, w\, \tfrac{1}{\kappa} \sum_k \Big[\bar{\psi}_{k\uparrow}\psi_{k\uparrow} + \bar{\psi}_{k\downarrow}\psi_{k\downarrow} \Big] \Big\} \times \\
&\qquad \exp\Big\{ \sum_l \Big[i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \bar{\xi}^l_{\uparrow\uparrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\uparrow}\psi_{k\uparrow} + i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \xi^l_{\uparrow\uparrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \bar{\psi}_{k\uparrow}\psi_{k\uparrow} \Big] \Big\} \times \\
&\qquad \exp\Big\{ \sum_l \Big[i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \bar{\xi}^l_{\downarrow\downarrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\downarrow}\psi_{k\downarrow} + i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \xi^l_{\downarrow\downarrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \bar{\psi}_{k\downarrow}\psi_{k\downarrow} \Big] \Big\} \times \\
&\qquad \exp\Big\{ \sum_l \Big[i\,\big(\tfrac{\lambda_l}{\kappa}\big)^{\frac{1}{2}}\, \bar{\xi}^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\uparrow}\psi_{k\downarrow} + i\,\big(\tfrac{\lambda_l}{\kappa}\big)^{\frac{1}{2}}\, \xi^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \bar{\psi}_{k\downarrow}\psi_{k\uparrow} \Big] \Big\} \times \\
&\qquad \exp\Big\{ \sum_l \Big[i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \bar{\phi}^l_{\uparrow\uparrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\uparrow}\bar{\psi}_{-k\uparrow} + i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \phi^l_{\uparrow\uparrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \psi_{k\uparrow}\psi_{-k\uparrow} \Big] \Big\} \times \\
&\qquad \exp\Big\{ \sum_l \Big[i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \bar{\phi}^l_{\downarrow\downarrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\downarrow}\bar{\psi}_{-k\downarrow} + i\,\big(\tfrac{\lambda_l}{2\kappa}\big)^{\frac{1}{2}}\, \phi^l_{\downarrow\downarrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \psi_{k\downarrow}\psi_{-k\downarrow} \Big] \Big\} \times \\
&\qquad \exp\Big\{ \sum_l \Big[i\,\big(\tfrac{\lambda_l}{\kappa}\big)^{\frac{1}{2}}\, \bar{\phi}^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\uparrow}\bar{\psi}_{-k\downarrow} + i\,\big(\tfrac{\lambda_l}{\kappa}\big)^{\frac{1}{2}}\, \phi^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \psi_{k\uparrow}\psi_{-k\downarrow} \Big] \Big\} d\nu(w,\xi,\phi)
\end{aligned}$$

where

$$d\nu(w,\xi,\phi) = \tfrac{1}{\sqrt{2\pi}}\, e^{-\frac{1}{2}w^2} dw \prod_l \prod_{\substack{(\sigma\tau)\in \\ \{\uparrow\uparrow,\downarrow\downarrow,\uparrow\downarrow\}}} \tfrac{1}{(2\pi)^2}\, e^{-\frac{1}{2}(|\phi^l_{\sigma\tau}|^2 + |\xi^l_{\sigma\tau}|^2)} du^l_{\sigma\tau} dv^l_{\sigma\tau} da^l_{\sigma\tau} db^l_{\sigma\tau}$$

By a substitution of variables and collecting terms, one obtains

$$\begin{aligned}
&e^{-\mathcal{U}(\psi,\bar{\psi})} = \\
&\int \exp\Big\{ \tfrac{1}{\kappa} \sum_k \Big[U_0^{\frac{1}{2}} w + i \sum_l \lambda_l^{\frac{1}{2}} \Big(\bar{\xi}^l_{\uparrow\uparrow}\, \bar{y}_l(\mathbf{k}') + \xi^l_{\uparrow\uparrow}\, y_l(\mathbf{k}') \Big) \Big] \bar{\psi}_{k\uparrow}\psi_{k\uparrow} \Big\} \times \\
&\quad \exp\Big\{ \tfrac{1}{\kappa} \sum_k \Big[U_0^{\frac{1}{2}} w + i \sum_l \lambda_l^{\frac{1}{2}} \Big(\bar{\xi}^l_{\downarrow\downarrow}\, \bar{y}_l(\mathbf{k}') + \xi^l_{\downarrow\downarrow}\, y_l(\mathbf{k}') \Big) \Big] \bar{\psi}_{k\downarrow}\psi_{k\downarrow} \Big\} \times \\
&\quad \exp\Big\{ \sum_l \Big[i(2\lambda_l)^{\frac{1}{2}} \bar{\xi}^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa} \sum_k \bar{y}_l(\mathbf{k}')\, \bar{\psi}_{k\uparrow}\psi_{k\downarrow} + i(2\lambda_l)^{\frac{1}{2}} \xi^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa} \sum_k y_l(\mathbf{k}')\, \bar{\psi}_{k\downarrow}\psi_{k\uparrow} \Big] \Big\} \times
\end{aligned}$$

$$
\begin{aligned}
&\exp\Big\{\sum_l \Big[i\lambda_l^{\frac12}\bar\phi^l_{\uparrow\uparrow}\, \tfrac{1}{\kappa}\sum_k \tfrac{\bar y_l(\mathbf{k}')-\bar y_l(-\mathbf{k}')}{2}\,\bar\psi_{k\uparrow}\bar\psi_{-k\uparrow} + i\lambda_l^{\frac12}\phi^l_{\uparrow\uparrow}\, \tfrac{1}{\kappa}\sum_k \tfrac{y_l(\mathbf{k}')-y_l(-\mathbf{k}')}{2}\,\psi_{k\uparrow}\psi_{-k\uparrow}\Big]\Big\}\times \\
&\exp\Big\{\sum_l \Big[i\lambda_l^{\frac12}\bar\phi^l_{\downarrow\downarrow}\, \tfrac{1}{\kappa}\sum_k \tfrac{\bar y_l(\mathbf{k}')-\bar y_l(-\mathbf{k}')}{2}\,\bar\psi_{k\downarrow}\bar\psi_{-k\downarrow} + i\lambda_l^{\frac12}\phi^l_{\downarrow\downarrow}\, \tfrac{1}{\kappa}\sum_k \tfrac{y_l(\mathbf{k}')-y_l(-\mathbf{k}')}{2}\,\psi_{k\downarrow}\psi_{-k\downarrow}\Big]\Big\}\times \\
&\exp\Big\{\sum_l \Big[i(2\lambda_l)^{\frac12}\bar\phi^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa}\sum_k \bar y_l(\mathbf{k}')\,\bar\psi_{k\uparrow}\bar\psi_{-k\downarrow} + i(2\lambda_l)^{\frac12}\phi^l_{\uparrow\downarrow}\, \tfrac{1}{\kappa}\sum_k y_l(\mathbf{k}')\,\psi_{k\uparrow}\psi_{-k\downarrow}\Big]\Big\} d\nu_\kappa(w,\xi,\phi)
\end{aligned}
$$

where $d\nu_\kappa(w,\xi,\phi)$ is defined in the statement of the theorem. Using the definition of the fields Ξ, Γ and Φ, the above expression reads

$$
\begin{aligned}
e^{-\mathcal{U}(\psi,\bar\psi)} = \int \exp\Big\{ \tfrac{1}{\kappa}\sum_k \Big[&\Gamma_{\mathbf{k}\uparrow}\,\bar\psi_{k\uparrow}\psi_{k\uparrow} + \Gamma_{\mathbf{k}\downarrow}\,\bar\psi_{k\downarrow}\psi_{k\downarrow} + i\bar\Xi_{\mathbf{k}\uparrow\downarrow}\bar\psi_{k\uparrow}\psi_{k\downarrow} + i\Xi_{\mathbf{k}\uparrow\downarrow}\bar\psi_{k\downarrow}\psi_{k\uparrow} \\
&+ i\bar\Phi_{\mathbf{k}\uparrow\downarrow}\bar\psi_{k\uparrow}\bar\psi_{-k\downarrow} + i\Phi_{\mathbf{k}\uparrow\downarrow}\psi_{k\uparrow}\psi_{-k\downarrow} + \tfrac{i}{2}\sum_\sigma \big(\bar\Phi_{\mathbf{k}\sigma\sigma}\bar\psi_{k\sigma}\bar\psi_{-k\sigma} + \Phi_{\mathbf{k}\sigma\sigma}\psi_{k\sigma}\psi_{-k\sigma}\big)\Big]\Big\} d\nu_\kappa
\end{aligned}
$$

We now rewrite the exponent in order to perform the fermionic functional integral. Since, if the set of spatial momenta satisfy $\mathbf{k} \in M_\omega = -M_\omega$,

$$
\begin{aligned}
\sum_k \Gamma_{k\uparrow}\,\bar\psi_{k\uparrow}\psi_{k\uparrow} &= \sum_{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)}\ \sum_{\mathbf{k}\in M_\omega} \Gamma_{k\uparrow}\,\bar\psi_{k\uparrow}\psi_{k\uparrow} = \sum_{k_0>0}\sum_{\mathbf{k}\in M_\omega}\Big[\Gamma_{k\uparrow}\,\bar\psi_{k\uparrow}\psi_{k\uparrow} + \Gamma_{-k\uparrow}\,\bar\psi_{-k\uparrow}\psi_{-k\uparrow}\Big] \\
&= \tfrac12 \sum_{\substack{k\\ k_0>0}} \Big[\Gamma_{k\uparrow}\big(\bar\psi_{k\uparrow}\psi_{k\uparrow} - \psi_{k\uparrow}\bar\psi_{k\uparrow}\big) + \Gamma_{-k\uparrow}\big(\bar\psi_{-k\uparrow}\psi_{-k\uparrow} - \psi_{-k\uparrow}\bar\psi_{-k\uparrow}\big)\Big]
\end{aligned}
$$

one obtains, using the antisymmetry of the $\Phi_{k\sigma\sigma}$, $\bar\Phi_{k\sigma\sigma}$

$$
\begin{aligned}
&\sum_k \Big\{ \textstyle\sum_\sigma \Gamma_{\mathbf{k}\sigma}\,\bar\psi_{k\sigma}\psi_{k\sigma} + i\bar\Phi_{\mathbf{k}\uparrow\downarrow}\bar\psi_{k\uparrow}\bar\psi_{-k\downarrow} + i\Phi_{\mathbf{k}\uparrow\downarrow}\psi_{k\uparrow}\psi_{-k\downarrow} \\
&\qquad + i\bar\Xi_{\mathbf{k}\uparrow\downarrow}\bar\psi_{k\uparrow}\psi_{k\downarrow} + i\Xi_{\mathbf{k}\uparrow\downarrow}\bar\psi_{k\downarrow}\psi_{k\uparrow} + \tfrac{i}{2}\textstyle\sum_\sigma\big(\bar\Phi_{\mathbf{k}\sigma\sigma}\bar\psi_{k\sigma}\bar\psi_{-k\sigma} + \Phi_{\mathbf{k}\sigma\sigma}\psi_{k\sigma}\psi_{-k\sigma}\big)\Big\} \\
&= \tfrac12 \sum_{\substack{k\\ k_0>0}} \Big\{ \textstyle\sum_\sigma \Big[\Gamma_{\mathbf{k}\sigma}\big(\bar\psi_{k\sigma}\psi_{k\sigma} - \psi_{k\sigma}\bar\psi_{k\sigma}\big) + \Gamma_{-\mathbf{k}\sigma}\big(\bar\psi_{-k\sigma}\psi_{-k\sigma} - \psi_{-k\sigma}\bar\psi_{-k\sigma}\big)\Big] \\
&\qquad + i\bar\Phi_{\mathbf{k}\uparrow\downarrow}\big(\bar\psi_{k\uparrow}\bar\psi_{-k\downarrow} - \bar\psi_{-k\downarrow}\bar\psi_{k\uparrow}\big) + i\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\big(\bar\psi_{-k\uparrow}\bar\psi_{k\downarrow} - \bar\psi_{k\downarrow}\bar\psi_{-k\uparrow}\big) \\
&\qquad + i\Phi_{\mathbf{k}\uparrow\downarrow}\big(\psi_{k\uparrow}\psi_{-k\downarrow} - \psi_{-k\downarrow}\psi_{k\uparrow}\big) + i\Phi_{-\mathbf{k}\uparrow\downarrow}\big(\psi_{-k\uparrow}\psi_{k\downarrow} - \psi_{k\downarrow}\psi_{-k\uparrow}\big) \\
&\qquad + i\bar\Xi_{\mathbf{k}\uparrow\downarrow}\big(\bar\psi_{k\uparrow}\psi_{k\downarrow} - \psi_{k\downarrow}\bar\psi_{k\uparrow}\big) + i\bar\Xi_{-\mathbf{k}\uparrow\downarrow}\big(\bar\psi_{-k\uparrow}\psi_{-k\downarrow} - \psi_{-k\downarrow}\bar\psi_{-k\uparrow}\big) \\
&\qquad + i\Xi_{\mathbf{k}\uparrow\downarrow}\big(\bar\psi_{k\downarrow}\psi_{k\uparrow} - \psi_{k\uparrow}\bar\psi_{k\downarrow}\big) + i\Xi_{-\mathbf{k}\uparrow\downarrow}\big(\bar\psi_{-k\downarrow}\psi_{-k\uparrow} - \psi_{-k\uparrow}\bar\psi_{-k\downarrow}\big) \\
&\qquad + i\textstyle\sum_\sigma\Big[\bar\Phi_{\mathbf{k}\sigma\sigma}\big(\bar\psi_{k\sigma}\bar\psi_{-k\sigma} - \bar\psi_{-k\sigma}\bar\psi_{k\sigma}\big) + \Phi_{\mathbf{k}\sigma\sigma}\big(\psi_{k\sigma}\psi_{-k\sigma} - \psi_{-k\sigma}\psi_{k\sigma}\big)\Big]\Big\}
\end{aligned}
$$

Since, if $a_k = ik_0 - e_{\mathbf{k}}$

$$
d\mu_C(\psi,\bar\psi) = \prod_{k,\sigma}\tfrac{\kappa}{a_k}\; e^{-\frac{1}{\kappa}\sum_{k,\sigma} a_k\bar\psi_{k,\sigma}\psi_{k,\sigma}} \prod_{k,\sigma} d\psi_{k,\sigma}d\bar\psi_{k,\sigma}
$$

one obtains

$$\begin{aligned}
Z(\beta, L, \{s_{k,\sigma}\}) &= \int e^{-\mathcal{U}(\psi,\bar\psi)+\frac{1}{\kappa}\sum_{k,\sigma} s_{k,\sigma}\bar\psi_{k,\sigma}\psi_{k,\sigma}}\,d\mu_C(\psi,\bar\psi) \\
&= \prod_{k,\sigma}\tfrac{\kappa}{a_k}\int\int e^{-\frac{1}{2}\frac{1}{\kappa}\sum_{k_0>0}\sum_{\mathbf{k}}\langle\Psi_k,S_k\Psi_k\rangle}\prod_{k,\sigma} d\psi_{k\sigma}d\bar\psi_{k\sigma}\,d\nu_\kappa(w,\xi,\phi) \\
&= \int\prod_{\substack{k_0>0\\ \mathbf{k}\in M_\omega}}\Big\{\Big(\tfrac{\kappa^2}{a_k a_{-k}}\Big)^2\int e^{-\frac{1}{2}\frac{1}{\kappa}\langle\Psi_k,S_k\Psi_k\rangle}d\bar\psi_{k\uparrow}d\psi_{k\uparrow}d\bar\psi_{-k\downarrow}d\psi_{-k\downarrow}\times \\
&\qquad\qquad d\psi_{k\downarrow}d\bar\psi_{k\downarrow}d\psi_{-k\uparrow}d\bar\psi_{-k\uparrow}\Big\}d\nu_\kappa(w,\xi,\phi) \\
&= \int\prod_{\substack{k_0>0\\ \mathbf{k}\in M_\omega}}\Big\{\Big(\tfrac{1}{a_k a_{-k}}\Big)^2\mathrm{Pf}S_k\Big\}d\nu_\kappa(w,\xi,\phi)
\end{aligned}$$

where

$$\Psi_k = \big(\bar\psi_{k\uparrow},\psi_{k\uparrow},\bar\psi_{-k\downarrow},\psi_{-k\downarrow},\psi_{k\downarrow},\bar\psi_{k\downarrow},\psi_{-k\uparrow},\bar\psi_{-k\uparrow}\big)$$

and $\mathrm{Pf}S_k$ is the Pfaffian of the 8×8 skew symmetric matrix S_k defined in the statement of the theorem. We used that

$$\prod_{k,\sigma} d\psi_{k\sigma}d\bar\psi_{k\sigma} = \prod_{k_0>0}\prod_{\mathbf{k}}\Big\{+d\bar\psi_{k\uparrow}d\psi_{k\uparrow}d\bar\psi_{-k\downarrow}d\psi_{-k\downarrow}d\psi_{k\downarrow}d\bar\psi_{k\downarrow}d\psi_{-k\uparrow}d\bar\psi_{-k\uparrow}\Big\}$$

Part b) of the theorem follows from

$$\tfrac{1}{\kappa}\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle = \tfrac{\partial}{\partial s_{p\sigma}}_{|s=0}\log Z(\beta,L,\{s_{k\sigma}\}) \qquad\bullet$$

Lemma II.2: *Let S_k be the skew symmetric 8×8 matrix of Theorem II.1. Then the Pfaffian of S_k is given by*

$$\begin{aligned}
\mathrm{Pf}S_k =&\big(A_{k\uparrow}A_{-k\downarrow}+\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\big)\big(A_{-k\uparrow}A_{k\downarrow}+\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\big) \\
&+A_{k\uparrow}A_{-k\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}+A_{k\downarrow}A_{-k\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow} \\
&+\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}+\bar\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} \\
&+\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow} \\
&+i\Xi_{\mathbf{k}\uparrow\downarrow}A_{-k\uparrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\downarrow\downarrow}+i\bar\Xi_{\mathbf{k}\uparrow\downarrow}A_{-k\uparrow}\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow} \\
&-i\Xi_{\mathbf{k}\uparrow\downarrow}A_{-k\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow}-i\bar\Xi_{\mathbf{k}\uparrow\downarrow}A_{-k\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow} \\
&+i\Xi_{-\mathbf{k}\uparrow\downarrow}A_{k\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow}+i\bar\Xi_{-\mathbf{k}\uparrow\downarrow}A_{k\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow} \\
&-i\Xi_{-\mathbf{k}\uparrow\downarrow}A_{k\uparrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\downarrow\downarrow}-i\bar\Xi_{-\mathbf{k}\uparrow\downarrow}A_{k\uparrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow} \\
&+\Xi_{\mathbf{k}\uparrow\downarrow}\bar\Xi_{\mathbf{k}\uparrow\downarrow}A_{-k\uparrow}A_{-k\downarrow}+\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Xi_{-\mathbf{k}\uparrow\downarrow}A_{k\uparrow}A_{k\downarrow} \\
&+\Xi_{\mathbf{k}\uparrow\downarrow}\bar\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}+\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Xi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow} \\
&-\Xi_{\mathbf{k}\uparrow\downarrow}\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\bar\Xi_{-\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow} \\
&+\Xi_{\mathbf{k}\uparrow\downarrow}\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Xi_{-\mathbf{k}\uparrow\downarrow}
\end{aligned}$$

Proof: The Pfaffian of S_k is given by the sum of all contractions $\sum\Pi\langle\psi\psi\rangle$ of the fields

$$\bar\psi_{k\uparrow},\psi_{k\uparrow},\bar\psi_{-k\downarrow},\psi_{-k\downarrow},\psi_{k\downarrow},\bar\psi_{k\downarrow},\psi_{-k\uparrow},\bar\psi_{-k\uparrow}$$

where the value $\langle\psi\psi\rangle$ is given by the corresponding matrix element. That is, the Pfaffian can be evaluated by using Wick's Theorem or integration by parts. Since S_k is an 8×8 matrix, one has $\mathrm{Pf}[-S_k]=\mathrm{Pf}S_k$ and

$$\begin{aligned}
\mathrm{Pf}S_k &=\langle\bar\psi_{k\uparrow}\psi_{k\uparrow}\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&=-A_{k\uparrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}-i\bar\Phi_{\mathbf{k}\uparrow\downarrow}\langle\psi_{k\uparrow}\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-i\bar\Xi_{\mathbf{k}\uparrow\downarrow}\langle\psi_{k\uparrow}\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}+i\bar\Phi_{\mathbf{k}\uparrow\uparrow}\langle\psi_{k\uparrow}\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\rangle_{S_k}\\
&=+A_{k\uparrow}A_{-k\downarrow}\langle\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}+iA_{k\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}\langle\psi_{-k\downarrow}\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad+iA_{k\uparrow}\Xi_{-\mathbf{k}\uparrow\downarrow}\langle\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad+\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\langle\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}-\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Xi_{\mathbf{k}\uparrow\downarrow}\langle\psi_{-k\downarrow}\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\langle\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Xi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\bar\psi_{k\downarrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad+\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\rangle_{S_k}-\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Xi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\uparrow\uparrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\rangle_{S_k}\\
&=+\big(A_{k\uparrow}A_{-k\downarrow}+\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\big)\langle\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad+\big(A_{k\uparrow}i\Xi_{-\mathbf{k}\uparrow\downarrow}-\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\big)\langle\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad+\big(A_{k\uparrow}i\bar\Phi_{\mathbf{k}\downarrow\downarrow}-\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Xi_{\mathbf{k}\uparrow\downarrow}\big)\langle\psi_{-k\downarrow}\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Xi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{-k\uparrow}\bar\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\bar\psi_{k\downarrow}\bar\psi_{-k\uparrow}\rangle_{S_k}+\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\psi_{-k\uparrow}\rangle_{S_k}\\
&\quad-\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Xi_{\mathbf{k}\uparrow\downarrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\psi_{-k\uparrow}\rangle_{S_k}-\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\uparrow\uparrow}\langle\bar\psi_{-k\downarrow}\psi_{-k\downarrow}\psi_{k\downarrow}\bar\psi_{k\downarrow}\rangle_{S_k}\\
&=+\big(A_{k\uparrow}A_{-k\downarrow}+\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\big)\big(A_{k\downarrow}A_{-k\uparrow}+\Phi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\big)\\
&\quad+\big(A_{k\uparrow}i\Xi_{-\mathbf{k}\uparrow\downarrow}-\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\big)\big(-\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}-i\bar\Xi_{-k\uparrow\downarrow}A_{k\downarrow}\big)\\
&\quad+\big(A_{k\uparrow}i\bar\Phi_{\mathbf{k}\downarrow\downarrow}-\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Xi_{\mathbf{k}\uparrow\downarrow}\big)\big(-i\Phi_{\mathbf{k}\downarrow\downarrow}A_{-k\uparrow}-\bar\Xi_{-k\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\Big)\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\big(-i\bar\Phi_{\mathbf{k}\downarrow\downarrow}A_{-k\uparrow}-\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow}\big)\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Xi_{\mathbf{k}\uparrow\downarrow}\big(-A_{-k\downarrow}A_{-k\uparrow}-\Xi_{-\mathbf{k}\uparrow\downarrow}\bar\Xi_{-\mathbf{k}\uparrow\downarrow}\big)\\
&\quad-\bar\Xi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\big(A_{-k\downarrow}i\bar\Phi_{-\mathbf{k}\uparrow\downarrow}+\bar\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Xi_{-k\uparrow\downarrow}\big)\\
&\quad+\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\uparrow\downarrow}\big(\bar\Phi_{\mathbf{k}\downarrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}+i\Xi_{-\mathbf{k}\uparrow\downarrow}A_{k\downarrow}\big)
\end{aligned}$$

$$
\begin{aligned}
&-\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Xi_{\mathbf{k}\uparrow\downarrow}\big(A_{-k\downarrow}i\Phi_{-\mathbf{k}\uparrow\downarrow}+\Xi_{-\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\downarrow\downarrow}\big)\\
&-\bar\Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\uparrow\uparrow}\big(-A_{-k\downarrow}A_{k\downarrow}-\bar\Phi_{\mathbf{k}\downarrow\downarrow}\Phi_{\mathbf{k}\downarrow\downarrow}\big)
\end{aligned}
$$

By multiplying out the brackets one obtains the stated formula •

Before we specialize Theorem II.1 in section III where the effective potential and the two point functions are computed more explicitly, in the following theorem we write down the integral representation for the generating functional of the connected amputated Greens functions.

Theorem II.3: *Let $\mathcal{U}(\psi,\bar\psi)$ be given by (II.1) and let V be the effective potential (II.14). Let*

$$
G(\eta)=\log\int e^{-\mathcal{U}(\psi+\eta,\bar\psi+\bar\eta)}d\mu_C(\psi,\bar\psi) \tag{II.16}
$$

be the generating functional for the connected amputated Greens functions. Then, if $\underline{\zeta}_k$ denotes the eight component vector

$$
\underline{\zeta}_k=(a_k\eta_{k\uparrow},-a_k\bar\eta_{k\uparrow},a_{-k}\eta_{-k\downarrow},-a_{-k}\bar\eta_{-k\downarrow},-a_k\bar\eta_{k\downarrow},a_k\eta_{k\downarrow},-a_{-k}\bar\eta_{-k\uparrow},a_{-k}\eta_{-k\uparrow})
$$

and $S_k=S_k(w,\xi,\phi)$ is the 8×8 matrix of Theorem II.1, one has the following integral representation

$$
G(\eta)-G(0)=-\tfrac{1}{\kappa}\sum_{k\sigma}a_k\bar\eta_{k\sigma}\eta_{k\sigma}+\log\frac{\int e^{-\frac12\frac1\kappa\sum_k\langle\underline{\zeta}_k,S_k^{-1}\underline{\zeta}_k\rangle}e^{-\kappa V(w,\xi,\phi)}dw\,\Pi\,d\xi d\phi}{\int\ e^{-\kappa V(w,\xi,\phi)}dw\,\Pi\,d\xi d\phi} \tag{II.17}
$$

Proof: By a substitution of Grassmann variables,

$$
\int e^{-\mathcal{U}(\psi+\eta,\bar\psi+\bar\eta)}d\mu_C=
$$

$$
e^{-\frac1\kappa\sum_{k\sigma}a_k\bar\eta_{k\sigma}\eta_{k\sigma}}\int e^{-\mathcal{U}(\psi,\bar\psi)}\prod_{k\sigma}\tfrac{\kappa}{a_k}\,e^{\frac1\kappa\sum_{k\sigma}a_k[\bar\psi_{k\sigma}\eta_{k\sigma}+\bar\eta_{k\sigma}\psi_{k\sigma}-\bar\psi_{k\sigma}\psi_{k\sigma}]}\prod_{k\sigma}d\psi_{k\sigma}d\bar\psi_{k\sigma}
$$

As in the proof of Theorem II.1, one has

$$
e^{-\mathcal{U}(\psi,\bar\psi)-\frac1\kappa\sum_{k\sigma}a_k\bar\psi_{k\sigma}\psi_{k\sigma}}=\int e^{-\frac12\frac1\kappa\sum_{k_0>0}\sum_{\mathbf{k}}\langle\Psi_k,S_k\Psi_k\rangle}d\nu_\kappa(w,\xi,\phi)
$$

such that

$$
\int e^{-\mathcal{U}(\psi,\bar\psi)}\prod_{k\sigma}\tfrac{\kappa}{a_k}\,e^{\frac1\kappa\sum_{k\sigma}a_k[\bar\psi_{k\sigma}\eta_{k\sigma}+\bar\eta_{k\sigma}\psi_{k\sigma}-\bar\psi_{k\sigma}\psi_{k\sigma}]}\prod_{k\sigma}d\psi_{k\sigma}d\bar\psi_{k\sigma}
$$

$$
\begin{aligned}
&= \int\int \prod_{k_0>0,\mathbf{k}} \left(\tfrac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2}\right) e^{\frac{1}{\kappa}\sum_{k\sigma} a_k[\bar\psi_{k\sigma}\eta_{k\sigma}+\bar\eta_{k\sigma}\psi_{k\sigma}]}\times \\
&\qquad \prod_{k_0>0,\mathbf{k}} \tfrac{\kappa^4}{\mathrm{Pf}S_k}\, e^{-\frac{1}{2}\frac{1}{\kappa}\sum_{k_0>0}\sum_{\mathbf{k}}\langle\Psi_k,S_k\Psi_k\rangle} d\nu_\kappa(w,\xi,\phi) \prod_{k\sigma} d\psi_{k\sigma} d\bar\psi_{k\sigma} \\
&= \int\int \prod_{k_0>0,\mathbf{k}} \left(\tfrac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2}\right) e^{\frac{1}{\kappa}\sum_{k_0>0,\mathbf{k}}\langle\Psi_k,\underline{\zeta}_k\rangle}\times \\
&\qquad \prod_{k_0>0,\mathbf{k}} \tfrac{\kappa^4}{\mathrm{Pf}S_k}\, e^{-\frac{1}{2}\frac{1}{\kappa}\sum_{k_0>0}\sum_{\mathbf{k}}\langle\Psi_k,S_k\Psi_k\rangle} \prod_{k_0>0,\mathbf{k}} d\Psi_k\, d\nu_\kappa(w,\xi,\phi) \\
&= \int \prod_{k_0>0,\mathbf{k}} \left(\tfrac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2}\right) e^{-\frac{1}{2}\frac{1}{\kappa}\sum_{k_0>0,\mathbf{k}}\langle\underline{\zeta}_k,S_k^{-1}\underline{\zeta}_k\rangle} d\nu_\kappa(w,\xi,\phi)
\end{aligned}
$$

where

$$
d\Psi_k = d\bar\psi_{k\uparrow} d\psi_{k\uparrow} d\bar\psi_{-k\downarrow} d\psi_{-k\downarrow} d\psi_{k\downarrow} d\bar\psi_{k\downarrow} d\psi_{-k\uparrow} d\bar\psi_{-k\uparrow}
$$

By definition of V

$$
\prod_{k_0>0,\mathbf{k}} \left(\tfrac{\mathrm{Pf}S_k}{a_k^2 a_{-k}^2}\right) d\nu_\kappa(w,\xi,\phi) = e^{-\kappa V(w,\xi,\phi)} dw \prod d\xi d\phi
$$

which proves the theorem •

III. Solution for Pure BCS

We consider the model

$$
Z(\beta, L, \{s_{k,\sigma}\}) = \int e^{-\mathcal{U}(\psi,\bar\psi)+\frac{1}{\kappa}\sum_{k,\sigma} s_{k,\sigma}\bar\psi_{k,\sigma}\psi_{k,\sigma}} d\mu_C(\psi,\bar\psi) \tag{III.1}
$$

with $e_{\mathbf{k}} = \mathbf{k}^2/2m - \mu$, $C(k) = 1/(ik_0 - e_{\mathbf{k}})$ and

$$
\mathcal{U}(\psi,\bar\psi) = \tfrac{1}{\kappa^3} \sum_{\sigma,\tau\in\{\uparrow,\downarrow\}} \sum_{k,p} U(\mathbf{k}-\mathbf{p})\, \bar\psi_{k,\sigma}\bar\psi_{-k,\tau}\psi_{p,\sigma}\psi_{-p,\tau} \tag{III.2}
$$

The electron-electron interaction U is given by (I.37).

To write down the effective potential and the two point functions in this case, one first has to compute the Pfaffian of the matrix S_k of Theorem II.1. Since we consider only a BCS interaction, the Ξ and Γ fields are zero. With Lemma II.2 one obtains

$$
\begin{aligned}
\mathrm{Pf}S_k =& \big(a_{k\uparrow}a_{-k\downarrow} + \bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\big)\big(a_{-k\uparrow}a_{k\downarrow} + \bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\big) \\
&+ a_{k\uparrow}a_{-k\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow} + a_{k\downarrow}a_{-k\downarrow}\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow} \\
&+ \Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow} + \bar\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} + \Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}
\end{aligned} \tag{III.3}
$$

where $a_{k\sigma} = a_k - s_{k\sigma}$. In particular, for $s_{k\sigma} = 0$

$$\mathrm{Pf} S_k = (a_k a_{-k} + \Omega^+_{\mathbf{k}})(a_k a_{-k} + \Omega^-_{\mathbf{k}}) \tag{III.4}$$

where $\Omega^\pm_{\mathbf{k}}$ are the solutions of the quadratic equation

$$\begin{aligned}\Omega^2 - \Big(\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow} + \bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} + \Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow} + \Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}\Big)\Omega + \Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow} \\ + \Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\downarrow}\bar\Phi_{-\mathbf{k}\uparrow\downarrow} + \bar\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow} + \Phi_{\mathbf{k}\uparrow\uparrow}\Phi_{\mathbf{k}\downarrow\downarrow}\bar\Phi_{\mathbf{k}\uparrow\uparrow}\bar\Phi_{\mathbf{k}\downarrow\downarrow} = 0\end{aligned} \tag{III.5}$$

The effective potential becomes

$$V(\phi) = \sum_{\sigma\tau\in\{\uparrow\uparrow,\downarrow\downarrow,\uparrow\downarrow\}} \sum_{l=0}^{N} |\phi^l_{\sigma\tau}|^2 - \tfrac{1}{\beta L^d} \sum_{\mathbf{k}\in M} \log \prod_{k_0>0} \Big\{ \tfrac{a_k a_{-k} + \Omega^+_{\mathbf{k}}}{a_k a_{-k}} \tfrac{a_k a_{-k} + \Omega^-_{\mathbf{k}}}{a_k a_{-k}} \Big\} \tag{III.6}$$

We start in section III.1 with the case of a delta function interaction, that is, $J = 0$ in (I.37). This reproduces the usual mean field results. In section III.2, we consider a more general electron-electron interaction of the form (I.37) with arbitrary J. One finds that the standard approach based on approximating the Hamiltonian by quadratic terms and imposing a self consistency equation may be misleading.

III.1 BCS with Delta Function Interaction

Corollary III.1: *Let $s_{k,\uparrow}, s_{k,\downarrow}, r_k, \bar r_k$ be some real or complex numbers, let*

$$\begin{aligned} Z(\beta, L, \{s_k\}, \{r_k\}) = \int \exp\Big\{ -\tfrac{\lambda}{\kappa^3} \textstyle\sum\limits_{k,p} \psi_{k\uparrow}\psi_{-k\downarrow}\bar\psi_{p\uparrow}\bar\psi_{-p\downarrow} + \tfrac{1}{\kappa}\textstyle\sum\limits_{k}[s_{k\uparrow}\bar\psi_{k\uparrow}\psi_{k\uparrow} \\ + s_{k\downarrow}\bar\psi_{k\downarrow}\psi_{k\downarrow} + r_k\psi_{k\uparrow}\psi_{-k\downarrow} - \bar r_k\bar\psi_{k\uparrow}\bar\psi_{-k\downarrow}] \Big\} d\mu_C(\psi,\bar\psi) \end{aligned} \tag{III.7}$$

and let

$$\tfrac{1}{\kappa}\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle_{\beta,L,r} = \tfrac{\partial}{\partial s_{p\sigma}} \log Z(\beta, L, \{s_k\}, \{r_k\})\big|_{s_k=0, r_k=r} \tag{III.8}$$

$$\tfrac{1}{\kappa}\langle\psi_{p\uparrow}\psi_{-p\downarrow}\rangle_{\beta,L,r} = \tfrac{\partial}{\partial r_p} \log Z(\beta, L, \{s_k\}, \{r_k\})\big|_{s_k=0, r_k=r} \tag{III.9}$$

Define the effective potential

$$V_{\beta,r}(u,v) = u^2 + (v + \tfrac{|r|}{\sqrt{\lambda}})^2 - \textstyle\int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, W_{\beta,\mathbf{k}}\big(\lambda(u^2+v^2)\big) \tag{III.10}$$

where

$$W_{\beta,\mathbf{k}}(y) = \tfrac{1}{\beta}\log\left[\tfrac{\cosh(\frac{\beta}{2}\sqrt{e^2_{\mathbf{k}}+y})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right]^2 \tag{III.11}$$

a) *Let* $a_p = ip_0 - e_{\mathbf{p}}$. *There are the two dimensional integral representations*

$$Z(\beta, L, r) = \tfrac{\kappa}{\pi} \int e^{-\kappa V_{\beta,r}(u,v)} dudv \tag{III.12}$$

$$\tfrac{1}{\kappa} \langle \bar{\psi}_{p,\sigma} \psi_{p,\sigma} \rangle_{\beta,L,r} = \frac{\int \frac{-a_{-p}}{a_p a_{-p} + \lambda(u^2+v^2)} \; e^{-\kappa V_{\beta,r}(u,v)} dudv}{\int e^{-\kappa V_{\beta,r}(u,v)} dudv} \tag{III.13}$$

$$\tfrac{1}{\kappa} \langle \psi_{p,\uparrow} \psi_{-p,\downarrow} \rangle_{\beta,L,r} = \frac{\int \frac{-i\sqrt{\lambda}\, e^{-i\alpha}(u-iv)}{a_p a_{-p} + \lambda(u^2+v^2)} \; e^{-\kappa V_{\beta,r}(u,v)} dudv}{\int e^{-\kappa V_{\beta,r}(u,v)} dudv}$$

$$\tfrac{1}{\kappa} \langle \bar{\psi}_{p,\uparrow} \bar{\psi}_{-p,\downarrow} \rangle_{\beta,L,r} = \frac{\int \frac{-i\sqrt{\lambda}\, e^{i\alpha}(u+iv)}{a_p a_{-p} + \lambda(u^2+v^2)} \; e^{-\kappa V_{\beta,r}(u,v)} dudv}{\int e^{-\kappa V_{\beta,r}(u,v)} dudv} \tag{III.14}$$

b) *Let* $V_\beta(\rho) := \rho^2 - \int \frac{d^d\mathbf{k}}{(2\pi)^d} W_{\beta,\mathbf{k}}(\lambda\rho^2)$ *and* $|\Delta|^2 = \lambda\rho_0^2$ *be given by* $V_\beta(\rho_0) = \min_{\rho \geq 0} V_\beta(\rho)$ *Then, if* $\Delta = \Delta(\lambda, \beta) = |\Delta|\, e^{-i\alpha}$, *one has*

$$\lim_{|r|\to 0} \lim_{L\to\infty} \tfrac{1}{\beta L^d} \log Z(\beta, L, r) = -V_\beta(|\Delta|) \tag{III.15}$$

$$\lim_{|r|\to 0} \lim_{L\to\infty} \tfrac{1}{\beta L^d} \langle \bar{\psi}_{p,\sigma} \psi_{p,\sigma} \rangle_{\beta,L,r} = \tfrac{-a_{-p}}{a_p a_{-p} + |\Delta|^2} \tag{III.16}$$

$$\lim_{|r|\to 0} \lim_{L\to\infty} \tfrac{1}{\beta L^d} \langle \psi_{p,\uparrow} \psi_{-p,\downarrow} \rangle_{\beta,L,r} = \tfrac{\Delta}{a_p a_{-p} + |\Delta|^2} \tag{III.17}$$

$$\lim_{|r|\to 0} \lim_{L\to\infty} \tfrac{1}{\beta L^d} \langle \bar{\psi}_{p,\uparrow} \bar{\psi}_{-p,\downarrow} \rangle_{\beta,L,r} = \tfrac{-\bar{\Delta}}{a_p a_{-p} + |\Delta|^2} \tag{III.18}$$

but

$$\lim_{L\to\infty} \lim_{|r|\to 0} \tfrac{1}{\kappa} \langle \psi_{p,\uparrow} \psi_{-p,\downarrow} \rangle_{\beta,L,r} = \lim_{L\to\infty} \lim_{|r|\to 0} \tfrac{1}{\kappa} \langle \bar{\psi}_{p,\uparrow} \psi_{-p,\downarrow} \rangle_{\beta,L,r} = 0 \tag{III.19}$$

Proof: a) For a δ function interaction one has $J = 0$ in Theorem II.1 and the fields (II.10) are zero, $\Phi_{\mathbf{k}\sigma\sigma} = 0$. The fields (II.9) become $\Phi_{\mathbf{k}\uparrow\downarrow} = (2\lambda)^{\frac{1}{2}} \phi^0_{\uparrow\downarrow} \equiv (2\lambda)^{\frac{1}{2}} \phi$. Since we added the $r\psi\psi$ and $\bar{r}\bar{\psi}\bar{\psi}$ terms to the exponent in (III.7) which were not present in Theorem II.1, these fields have to be substituted by ($g = \sqrt{2\lambda}$)

$$g\phi \to g\phi - ir_k\,, \quad g\bar{\phi} \to g\bar{\phi} + i\bar{r}_k$$

The Pfaffian (III.3) becomes

$$\mathrm{Pf} S_k = \Big(a_{k\uparrow} a_{-k\downarrow} + [g\bar{\phi} + i\bar{r}_k][g\phi - ir_k] \Big) \Big(a_{-k\uparrow} a_{k\downarrow} + [g\bar{\phi} + i\bar{r}_{-k}][g\phi - ir_{-k}] \Big) \tag{III.20}$$

such that

$$\frac{\frac{\partial}{\partial s_{p\sigma}}\mathrm{Pf}S_p}{\mathrm{Pf}S_p}\Big|_{s_k=0,r_k=r} = \frac{-a_{-p}}{a_pa_{-p}+[g\phi-ir][g\bar{\phi}+i\bar{r}]} \tag{III.21}$$

and

$$\frac{\frac{\partial}{\partial r_p}\mathrm{Pf}S_p}{\mathrm{Pf}S_p}\Big|_{s_k=0,r_k=r} = \frac{-i[g\bar{\phi}+i\bar{r}]}{a_pa_{-p}+[g\phi-ir][g\bar{\phi}+i\bar{r}]} \tag{III.22}$$

The effective potential (III.6) is given by (the $|\phi|^2$ term is not shifted by r)

$$V(\phi) = |\phi|^2 - \frac{1}{\beta L^d}\sum_{\mathbf{k}\in M}\log\prod_{k_0>0}\left\{\frac{a_ka_{-k}+[g\phi-ir][g\bar{\phi}+i\bar{r}]}{a_ka_{-k}}\right\}^2 \tag{III.23}$$

The product over $k_0 \in \frac{\pi}{\beta}(2\mathbf{N}+1)$ in (III.23) can be computed explicitly using the formula [Hn]

$$\prod_{n=0}^{\infty}\left(1+\frac{\xi}{(2n+1)^2+a^2}\right) = \frac{\cosh\left(\frac{\pi}{2}\sqrt{a^2+\xi}\right)}{\cosh\frac{\pi}{2}a} \tag{III.24}$$

One obtains

$$\prod_{k_0>0}\left\{\frac{a_ka_{-k}+\xi}{a_ka_{-k}}\right\}^2 = \prod_{k_0\in\frac{\pi}{\beta}(2\mathbf{N}+1)}\left(1+\frac{\xi}{k_0^2+e_{\mathbf{k}}^2}\right)^2 = \left\{\frac{\cosh\left(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\xi}\right)}{\cosh\left(\frac{\beta}{2}e_{\mathbf{k}}\right)}\right\}^2 \tag{III.25}$$

which gives, if one approximates the Riemannian sum $\frac{1}{L^d}\sum_{\mathbf{k}\in M_\omega}$ by an integral

$$V(\phi) = |\phi|^2 - \int_{M_\omega}\frac{d^d\mathbf{k}}{(2\pi)^d}\,W_{\beta,\mathbf{k}}\big([g\phi-ir][g\bar{\phi}+i\bar{r}]\big) \tag{III.26}$$

Thus one arrives at the integral representations

$$Z(\beta,L,r) = \frac{\kappa}{\pi}\int e^{-\kappa V(\phi)}dudv \tag{III.27}$$

$$\frac{1}{\kappa}\langle\bar{\psi}_{p,\sigma}\psi_{p,\sigma}\rangle_{\beta,L,r} = \frac{\int\frac{-a_{-p}}{a_pa_{-p}+[g\phi-ir][g\bar{\phi}+i\bar{r}]}\,e^{-\kappa V(\phi)}dudv}{\int e^{-\kappa V(\phi)}dudv} \tag{III.28}$$

$$\frac{1}{\kappa}\langle\psi_{p,\uparrow}\psi_{-p,\downarrow}\rangle_{\beta,L,r} = \frac{\int\frac{-i[g\bar{\phi}+i\bar{r}]}{a_pa_{-p}+[g\phi-ir][g\bar{\phi}+i\bar{r}]}\,e^{-\kappa V(\phi)}dudv}{\int e^{-\kappa V(\phi)}dudv} \tag{III.29}$$

where V is given by (III.26). Part (a) then follows from the substitution of variables

$$\int F\left(g\phi-ir, g\bar{\phi}+i\bar{r}\right)\,e^{-\kappa|\phi|^2}dudv = \int F\left(e^{i\alpha}g\phi, e^{-i\alpha}g\bar{\phi}\right)e^{-\kappa\left(u^2+(v+\frac{|r|}{g})^2\right)}dudv \tag{III.30}$$

which holds for both signs of λ.

To obtain part **(b)**, one has to compute the limit of

$$\delta_{\kappa,r}(u,v) = \frac{e^{-\kappa V_{\beta,r}(u,v)}}{\int e^{-\kappa V_{\beta,r}(u,v)}dudv} \tag{III.31}$$

where

$$V_{\beta,r}(u,v) = u^2 + (v + \tfrac{|r|}{g})^2 - W_\beta\big(\lambda(u^2+v^2)\big) \tag{III.32}$$

if we define

$$W_\beta(\xi) = \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, W_{\beta,\mathbf{k}}(\xi) = \tfrac{1}{\beta}\int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\ \log\left[\frac{\cosh\big(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\xi}\big)}{\cosh\big(\frac{\beta}{2}e_{\mathbf{k}}\big)}\right]^2 \tag{III.33}$$

In particular,

$$\text{sign}W_\beta\big(\lambda(u^2+v^2)\big) = \text{sign}\lambda$$

For positive λ and nonzero r, $V_{\beta,r}(u,v)$ is real and has a unique global minimum determined by

$$\begin{aligned} 2u - 2u\lambda W_\beta'\big(\lambda(u^2+v^2)\big) &= 0 \\ 2(v+|r|) - 2v\lambda W_\beta'\big(\lambda(u^2+v^2)\big) &= 0 \end{aligned}$$

Since $\lambda W_\beta' = 1$ does not solve the second equation, the only solution of the first equation is $u = 0$ and one is left with

$$v\left[\lambda W_\beta'(\lambda v^2) - 1\right] = |r|$$

which has a solution $v = O(|r|)$ which is a local maximum and two nontrivial solutions $\lambda v^2 = |\Delta|^2 + O(|r|)$ where the positive one is only a local minimum and the negative one, v_0, is the global minimum. Therefore

$$\lim_{\kappa\to\infty} \delta_{\kappa,r}(u,v) = \lim_{\kappa\to\infty} \frac{e^{-\kappa[V_{\beta,r}(u,v)-V_{\beta,r}(0,v_0)]}}{\int e^{-\kappa[V_{\beta,r}(u,v)-V_{\beta,r}(0,v_0)]}dudv} = \delta(u)\,\delta(v-v_0)$$

and

$$\lim_{|r|\to 0}\lim_{\kappa\to\infty} \delta_{\kappa,r}(u,v) = \delta(u)\,\delta\left(v + \tfrac{|\Delta|}{g}\right) \tag{III.34}$$

This proves the formulae under (b) for attractive λ. Since

$$\lim_{|r|\to 0} V_{r,\beta}(u,v) = u^2 + v^2 - W_\beta\left(\lambda(u^2+v^2)\right)$$

is an even function in u and v, $\lim_{r\to 0}\langle\psi_{p,\uparrow}\psi_{-p,\downarrow}\rangle_{\beta,L,r}$=$\lim_{r\to 0}\langle\bar\psi_{p,\uparrow}\bar\psi_{-p,\downarrow}\rangle_{\beta,L,r} = 0$. The limit of the logarithm of the partition function becomes

$$\begin{aligned} \tfrac{1}{\kappa}\log Z &= \tfrac{1}{\kappa}\log\tfrac{\kappa}{\pi}\int e^{-\kappa V_{\beta,r}(u,v)}dudv \\ &= \tfrac{1}{\kappa}\log\tfrac{\kappa}{\pi}\int e^{-\kappa[V_{\beta,r}(u,v)-V_{\beta,r}(0,v_0)]}dudv + \tfrac{1}{\kappa}\log e^{-\kappa V_{\beta,r}(0,v_0)} \end{aligned}$$

The first term on the right hand side may be approximated by $(V_{uv} = \frac{\partial^2 V_{\beta,r}}{\partial u \partial v}(0, v_0) = 0)$

$$\tfrac{1}{\kappa} \log \tfrac{\kappa}{\pi} \int e^{-\frac{\kappa}{2}(V_{uu}u^2 + V_{vv}(v-v_0)^2)} du dv = \tfrac{1}{\kappa} \log \tfrac{1}{\pi} \int e^{-\frac{1}{2}(V_{uu}u^2 + V_{vv}v^2)} du dv \overset{\kappa \to \infty}{\to} 0$$

which results in

$$\lim_{\kappa \to \infty} \tfrac{1}{\kappa} \log Z = -V_{\beta,r}(0, v_0)$$

Now let λ be negative. In that case the effective potential (III.32) is complex:

$$\begin{aligned} V_{\beta,r}(u,v) &= u^2 + v^2 - \tfrac{|r|^2}{|\lambda|} - 2iv \tfrac{|r|}{\sqrt{|\lambda|}} - W_\beta\left(\lambda(u^2+v^2)\right) \\ &= \rho^2 + |W_\beta\left(\lambda\rho^2\right)| - \tfrac{|r|^2}{|\lambda|} - 2i\rho\cos\varphi \tfrac{|r|}{\sqrt{|\lambda|}} \end{aligned}$$

Since the real part $U_{\beta,r} = \mathrm{Re} V_{\beta,r}$ has a global minimum at $u = v = 0$ one has, since $U'' = \frac{\partial^2 U_{\beta,r}}{\partial u^2}(0,0) = \frac{\partial^2 U_{\beta,r}}{\partial v^2}(0,0) > 0$,

$$\begin{aligned} \delta_{\kappa,r}(u,v) &= \frac{e^{-\kappa\left[U_{\beta,r}(u,v) - 2iv\frac{|r|}{\sqrt{|\lambda|}}\right]}}{\int e^{-\kappa\left[U_{\beta,r}(u,v) - 2iv\frac{|r|}{\sqrt{|\lambda|}}\right]} du dv} \approx \frac{e^{-\kappa\left[\frac{U''}{2}(u^2+v^2) - 2iv\frac{|r|}{\sqrt{|\lambda|}}\right]}}{\int e^{-\kappa\left[\frac{U''}{2}(u^2+v^2) - 2iv\frac{|r|}{\sqrt{|\lambda|}}\right]} du dv} \\ &= \frac{e^{-\kappa\frac{U''}{2}u^2}}{\int e^{-\kappa\frac{U''}{2}u^2} du} \; \frac{e^{-\kappa\left[\frac{U''}{2}v^2 - 2iv\frac{|r|}{\sqrt{|\lambda|}}\right]}}{\int e^{-\kappa\left[\frac{U''}{2}v^2 - 2iv\frac{|r|}{\sqrt{|\lambda|}}\right]} dv} \\ &= \frac{e^{-\kappa\frac{U''}{2}u^2}}{\int e^{-\kappa\frac{U''}{2}u^2} du} \; \frac{e^{-\kappa\frac{U''}{2}\left[v - i\frac{2|r|}{\sqrt{|\lambda|}U''}\right]^2}}{\int e^{-\kappa\frac{U''}{2}\left[v - i\frac{2|r|}{\sqrt{|\lambda|}U''}\right]^2} dv} \overset{\kappa\to\infty}{\to} \delta(u)\, \delta\Big(v - i\frac{2|r|}{\sqrt{|\lambda|}U''}\Big) \end{aligned}$$

which results in

$$\lim_{|r|\to 0} \lim_{\kappa\to\infty} \delta_{\kappa,r}(u,v) = \delta(u)\, \delta(v) \tag{III.35}$$

and

$$\lim_{|r|\to 0} \lim_{\kappa\to\infty} \tfrac{1}{\kappa} \langle \bar\psi_{p,\uparrow} \psi_{-p,\downarrow} \rangle_{\beta,L,r} = \int \tfrac{-a_{-p}}{a_p a_{-p} + \lambda(u^2+v^2)}\, \delta(u)\delta(v)\, du dv = \tfrac{-a_{-p}}{a_p a_{-p}} = -\tfrac{1}{a_p}$$

$$\lim_{|r|\to 0} \lim_{\kappa\to\infty} \tfrac{1}{\kappa} \langle \psi_{p,\uparrow} \psi_{-p,\downarrow} \rangle_{\beta,L,r} = \lim_{|r|\to 0} \lim_{\kappa\to\infty} \tfrac{1}{\kappa} \langle \bar\psi_{p,\uparrow} \bar\psi_{-p,\downarrow} \rangle_{\beta,L,r} = 0 \qquad \bullet$$

Using (II.17) and (III.34,35) the infinite volume limit for the generating functional for the connected Greens functions can be computed in a similar way. One finds

Corollary III.2: *Let* $r = |r|e^{i\alpha}$, $\gamma = ge^{i\alpha}(u+iv)$, $\bar\gamma = ge^{-i\alpha}(u-iv)$ *and let*

$$G(\eta) = G_r(\beta, L, \eta) = \log \int e^{\mathcal{U}_r(\psi+\eta, \bar\psi+\bar\eta)} d\mu_C(\psi, \bar\psi) \tag{III.36}$$

be the generating functional for the connected amputated Greens functions where

$$\mathcal{U}_r(\psi, \bar\psi) = -\tfrac{\lambda}{\kappa^3} \sum_{k,p} \psi_{k\uparrow}\psi_{-k\downarrow}\bar\psi_{p\uparrow}\bar\psi_{-p\downarrow} + \tfrac{1}{\kappa} \sum_k [r\psi_{k\uparrow}\psi_{-k\downarrow} - \bar r \bar\psi_{k\uparrow}\bar\psi_{-k\downarrow}] \tag{III.37}$$

Let $V_{\beta,r}$ *be the effective potential (III.10) and let* Δ *be given as in Corollary III.1. Then there is the integral representation*

$$\begin{aligned} G(\eta) - G(0) = &-\tfrac{1}{\kappa} \sum_k \Big[a_k \bar\eta_{k\uparrow}\eta_{k\uparrow} + a_{-k}\bar\eta_{-k\downarrow}\eta_{-k\downarrow}\Big] + \\ &+ \log \frac{\int dudv\, e^{-\kappa V_{\beta,r}(u,v)} \exp\Big\{ \frac{1}{\kappa}\sum_k (a_k\bar\eta_{k\uparrow}, -a_{-k}\eta_{-k\downarrow}) \frac{1}{a_k a_{-k} + \gamma\bar\gamma} \begin{pmatrix} a_{-k} & -i\bar\gamma \\ i\gamma & -a_k \end{pmatrix} \begin{pmatrix} a_k\eta_{k\uparrow} \\ -a_{-k}\bar\eta_{-k\downarrow} \end{pmatrix} \Big\}}{\int dudv\, e^{-\kappa V_{\beta,r}(u,v)}} \end{aligned} \tag{III.38}$$

For attractive $\lambda > 0$ *the infinite volume limit of the generating functional is given by*

$$\begin{aligned} \lim_{|r|\to 0} \lim_{L\to\infty} \{G(\beta, L, r, \eta) - G(\beta, L, r, 0)\} = &\\ -\tfrac{1}{\kappa} \sum_k \Big[a_k \tfrac{|\Delta|^2}{a_k a_{-k} + |\Delta|^2}\, \bar\eta_{k\uparrow}\eta_{k\uparrow} &+ a_{-k} \tfrac{|\Delta|^2}{a_k a_{-k} + |\Delta|^2}\, \bar\eta_{-k\downarrow}\eta_{-k\downarrow} \\ &+ \Delta \tfrac{a_k a_{-k}}{a_k a_{-k} + |\Delta|^2}\, \eta_{-k\downarrow}\eta_{k\uparrow} + \bar\Delta \tfrac{a_k a_{-k}}{a_k a_{-k} + |\Delta|^2}\, \bar\eta_{k\uparrow}\bar\eta_{-k\downarrow} \Big] \end{aligned} \tag{III.39}$$

For repulsive $\lambda < 0$ *one obtains*

$$\lim_{|r|\to 0} \lim_{L\to\infty} \{G(\beta, L, r, \eta) - G(\beta, L, r, 0)\} = 0\,. \tag{III.40}$$

III.2 BCS with Higher Angular Momentum Terms

We now consider the case where the electron-electron interaction contains higher angular momentum terms. In this case one finds that the usual mean field approach, based on approximating the Hamiltonian by quadratic terms, may be misleading. To simplify the algebra we assume that only even ℓ terms contribute in (I.37). In that case the fields $\Phi_{\mathbf{k}\sigma\sigma}$ with equal spin (II.10) are still zero. So let

$$U(\mathbf{k}'-\mathbf{p}') = \begin{cases} \frac{1}{2}\sum\limits_{\substack{\ell=-j \\ \ell \text{ even}}}^{j} \lambda_{|\ell|} e^{i\ell\varphi_{\mathbf{k}}} e^{-i\ell\varphi_{\mathbf{p}}} + \frac{\lambda_0}{2} & \text{if } d=2 \\ \sum\limits_{\substack{\ell=0 \\ \ell \text{ even}}}^{j} \sum\limits_{m=-\ell}^{\ell} \lambda_\ell \bar{Y}_{\ell m}(\mathbf{k}')\, Y_{\ell m}(\mathbf{p}') & \text{if } d=3 \end{cases} =: \sum_{\substack{l=0 \\ l \text{ even}}}^{J} \lambda_l\, y_l(\mathbf{k}')\, \bar{y}_l(\mathbf{p}') \tag{III.41}$$

where $y_l(-\mathbf{k}) = (-1)^l y_l(\mathbf{k})$. We only consider the $\bar\psi\psi$ expectations, so we let $r=0$. The Pfaffian (III.3) becomes

$$\mathrm{Pf} S_k = \big(a_{k\uparrow}a_{-k\downarrow} + \bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\big)\big(a_{-k\uparrow}a_{k\downarrow} + \bar\Phi_{-\mathbf{k}\uparrow\downarrow}\Phi_{-\mathbf{k}\uparrow\downarrow}\big) \tag{III.42}$$

where

$$\Phi_{\mathbf{k}\uparrow\downarrow} = \sum_{\substack{l=0 \\ l \text{ even}}}^{J} (2\lambda_l)^{\frac{1}{2}} \phi^l_{\uparrow\downarrow}\, y_l(\mathbf{k}')\,, \quad \bar\Phi_{\mathbf{k}\uparrow\downarrow} = \sum_{\substack{l=0 \\ l \text{ even}}}^{J} (2\lambda_l)^{\frac{1}{2}} \bar\phi^l_{\uparrow\downarrow}\, \bar{y}_l(\mathbf{k}') \tag{III.43}$$

and the effective potential is given by, using (III.25) again to compute the k_0 product

$$V(\phi_{\uparrow\downarrow}) = \sum_{l=0}^{J} |\phi^l_{\uparrow\downarrow}|^2 - \int_{M_\omega} \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, W_{\beta,\mathbf{k}}\big(\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}\big) \tag{III.44}$$

where $W_{\beta,\mathbf{k}}$ is given by (III.11). By Theorem II.1, the two point function is given by

$$\tfrac{1}{\kappa}\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle = -\frac{\int \frac{ip_0+e_{\mathbf{p}}}{p_0^2+e_{\mathbf{p}}^2+\bar\Phi_{\mathbf{k}\uparrow\downarrow}\Phi_{\mathbf{k}\uparrow\downarrow}}\, e^{-\kappa V(\phi_{\uparrow\downarrow})} \prod\limits_{l=0}^{J} du^l_{\uparrow\downarrow} dv^l_{\uparrow\downarrow}}{\int e^{-\kappa V(\phi_{\uparrow\downarrow})} \prod\limits_{l=0}^{J} du^l_{\uparrow\downarrow} dv^l_{\uparrow\downarrow}} \tag{III.45}$$

To compute the infinite volume limit, one has to find the global minimum of the real part of $V(\phi_{\uparrow\downarrow})$. This is easier for $d=2$. However, using symmetry arguments, it is possible to give a rather explicit expression also for $d=3$ without knowing the exact location of the global minimum.

Consider first the two dimensional case. An analysis done by Albrecht Schuette in his Diploma thesis [Sch] shows the following result:

Suppose that $\lambda_m > 0$ is attractive and $\lambda_m > \lambda_\ell$ for all $\ell \neq m$. Then

$$\lim_{\kappa\to\infty} \tfrac{1}{\kappa}\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle_{\beta,L} = \tfrac{ip_0+e_{\mathbf{p}}}{p_0^2+e_{\mathbf{p}}^2+\lambda_m\rho_m^2} \tag{III.46}$$

where ρ_m is determined by the BCS equation

$$1 - \tfrac{\lambda_m}{4\pi}\int d|\mathbf{k}||\mathbf{k}| \frac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\lambda_m\rho_m^2})}{\sqrt{e_{\mathbf{k}}^2+\lambda\rho_m^2}} = 0 \tag{III.47}$$

The form of the two point function (III.46) can still be obtained by applying the standard mean field formalism [AB,BW]. However, the situation is different in 3 dimensions. Before we state the corresponding theorem, we shortly recall the mean field equations (I.57,58)

The $\langle a^+ a\rangle$ expectations are given by

$$\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}\rangle = \tfrac{1}{2}\left(1 - e_{\mathbf{k}}\left[\tfrac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}})}{\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}}}\right]_{\sigma\sigma}\right) \tag{I.57}$$

where the 2×2 matrix $\Delta_{\mathbf{k}}$, $\Delta^T_{\mathbf{k}} = -\Delta_{-\mathbf{k}}$, is a solution of the gap equation

$$\Delta_{\mathbf{p}} = \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, U(\mathbf{p'}-\mathbf{k'})\, \Delta_{\mathbf{k}} \tfrac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}})}{2\sqrt{e_{\mathbf{k}}^2+\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}}}} \tag{I.58}$$

In 2 dimensions, if one substitutes $U(\mathbf{p}-\mathbf{k})$ by a single attractive term $\lambda_\ell e^{i\ell(\varphi_{\mathbf{p}}-\varphi_{\mathbf{k}})}$, then (I.58) has the unitary isotropic solution $\Delta_{\mathbf{k}} = \Delta\left(\begin{smallmatrix} 0 & e^{i\ell\varphi_{\mathbf{k}}} \\ -e^{i\ell\varphi_{\mathbf{k}}} & 0\end{smallmatrix}\right)$ for even ℓ and $\Delta_{\mathbf{k}} = \Delta\left(\begin{smallmatrix} \cos\ell\varphi_{\mathbf{k}} & \sin\ell\varphi_{\mathbf{k}} \\ \sin\ell\varphi_{\mathbf{k}} & -\cos\ell\varphi_{\mathbf{k}}\end{smallmatrix}\right)$ if ℓ is odd. In that case $\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}} = |\Delta|^2 Id$ and (I.57) coincides with (III.46) (after integration of the latter over p_0).

In 3 dimensions, it is proven [FKT2] that for all $\ell \geq 2$ (I.58) does not have unitary isotropic ($\Delta^*_{\mathbf{k}}\Delta_{\mathbf{k}} = const\, Id$) solutions. In view of that result, the symmetry considerations below indicate that in 3 dimensions for $\ell \geq 2$ the standard mean field approach is misleading since one would no longer expect $SO(3)$ invariance for the $\langle a^+_{\mathbf{k}\sigma} a_{\mathbf{k}\sigma}\rangle$ expectations according to (I.57). But this is indeed the case if there is no external SO(3) symmetry breaking term (which is also not present in the quadratic mean field model, see the discussion following (I.61) at the end of the introduction).

Theorem III.3: *Let ℓ be even, $\lambda_\ell > 0$ be attractive and let*

$$\mathcal{U}(\psi,\bar\psi) = \tfrac{\lambda_\ell}{(\beta L^d)^3} \sum_{k,p}\sum_{m=-\ell}^{\ell} \bar Y_{\ell m}(\mathbf{k'}) Y_{\ell m}(\mathbf{p'})\, \bar\psi_{k\uparrow}\bar\psi_{-k\downarrow}\psi_{p\uparrow}\psi_{-p\downarrow} \tag{III.48}$$

Then, if $e_{R\mathbf{k}} = e_{\mathbf{k}}$ for all $R \in SO(3)$, one has

$$\begin{aligned}\lim_{\kappa\to\infty} \tfrac{1}{\kappa}\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle_{\beta,L} &= \lim_{\kappa\to\infty}\int \tfrac{1}{\kappa}\bar\psi_{p\sigma}\psi_{p\sigma}\, e^{-\mathcal{U}(\psi,\bar\psi)} d\mu_C(\psi,\bar\psi) \\ &= \int_{S^2} \tfrac{ip_0+e_{\mathbf{p}}}{p_0^2+e_{\mathbf{p}}^2+\lambda_\ell\rho_0^2|\,\Sigma_m\,\alpha^0_m Y_{\ell m}(\mathbf{x})|^2}\, \tfrac{d\Omega(\mathbf{x})}{4\pi} \end{aligned}\tag{III.49}$$

where $\rho_0 \geq 0$ and $\alpha^0 \in \mathbf{C}^{2\ell+1}$, $\Sigma_m |\alpha^0_m|^2 = 1$, are values at the global minimum (which is degenerated) of

$$V(\rho,\alpha) = \rho^2 - \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, \tfrac{1}{\beta}\log\left[\tfrac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\lambda_\ell\rho^2|\,\Sigma_m\,\alpha_m Y_{\ell m}(\mathbf{k'})|^2})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right]^2$$

In particular, the momentum distribution $n_{\mathbf{p}}$ *is given by*

$$n_{\mathbf{p}} = \lim_{L\to\infty} \tfrac{1}{L^d} \langle a^+_{\mathbf{p}\sigma} a_{\mathbf{p}\sigma}\rangle_{\beta,L} = \int_{S^2} \tfrac{1}{2}\left(1 - e_{\mathbf{p}} \tfrac{\tanh(\frac{\beta}{2}\sqrt{e_{\mathbf{p}}^2+|\Delta(\mathbf{x})|^2})}{\sqrt{e_{\mathbf{p}}^2+|\Delta(\mathbf{x})|^2}}\right) \tfrac{d\Omega(\mathbf{x})}{4\pi} \tag{III.50}$$

and has $SO(3)$ *symmetry. Here* $\Delta(\mathbf{x}) = \lambda_\ell^{\frac{1}{2}} \rho_0 \sum_m \alpha^0_m Y_{\ell m}(\mathbf{x})$.

Proof: Substituting $u^l_{\uparrow\downarrow}$ by u_m and $u^l_{\uparrow\downarrow}$ by $-v_m$ in (III.45), one has to compute the infinite volume limit of

$$\tfrac{1}{\kappa}\langle\bar\psi_{p\sigma}\psi_{p\sigma}\rangle = -\frac{\int_{\mathbf{R}^{4\ell+2}} \frac{a_{-p}}{a_p a_{-p}+|\Phi_{\mathbf{p}}|^2}\, e^{-\kappa V(\phi)} \prod_{m=-\ell}^{\ell} du_m dv_m}{\int_{\mathbf{R}^{4\ell+2}} e^{-\kappa V(\phi)} \prod_{m=-\ell}^{\ell} du_m dv_m} \tag{III.51}$$

where

$$V(\phi) = \sum_{m=-\ell}^{\ell} |\phi_m|^2 - \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, \tfrac{2}{\beta} \log\left[\tfrac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+|\Phi_{\mathbf{k}}|^2})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right]$$

and

$$\Phi_{\mathbf{k}} = \lambda_\ell^{\frac{1}{2}} \sum_{m=-\ell}^{\ell} \bar\phi_m Y_{\ell m}(\mathbf{k}')$$

Let $U(R)$ be the unitary representation of $SO(3)$ given by

$$Y_{\ell m}(R\mathbf{k}') = \sum_{m'} U(R)_{mm'} Y_{\ell m'}(\mathbf{k}')$$

and let $\sum_m (\overline{U\phi})_m Y_{\ell m}(\mathbf{k}') =: (U\Phi)_{\mathbf{k}}$. Then for all $R\in SO(3)$ one has $\big(U(R)\Phi\big)_{\mathbf{k}} = \Phi_{R^{-1}\mathbf{k}}$ and

$$\begin{aligned} V\big(U(R)\phi\big) &= \sum_m |[U(R)\phi]_m|^2 - \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, \tfrac{2}{\beta} \log\left[\tfrac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+|[U\Phi]_{\mathbf{k}}|^2})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right] \\ &= \sum_m |\phi_m|^2 - \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, \tfrac{2}{\beta} \log\left[\tfrac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+|\Phi_{R^{-1}\mathbf{k}}|^2})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right] \\ &= \sum_m |\phi_m|^2 - \int_M \tfrac{d^d\mathbf{k}}{(2\pi)^d}\, \tfrac{2}{\beta} \log\left[\tfrac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+|\Phi_{\mathbf{k}}|^2})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right] \;=\; V(\phi) \end{aligned} \tag{III.52}$$

Let $S^{4\ell+1} = \{\phi\in\mathbf{C}^{2\ell+1} \mid \sum_m|\phi_m|^2 = 1\}$. Since $U(R)$ leaves $S^{4\ell+1}$ invariant, $S^{4\ell+1}$ can be written as the union of disjoint orbits,

$$S^{4\ell+1} = \cup_{[\alpha]\in O}[\alpha]$$

where $[\alpha] = \{U(R)\alpha \mid R\in SO(3)\}$ is the orbit of $\alpha\in S^{4\ell+1}$ under the action of $U(R)$ and O is the set of all orbits. If one chooses a fixed representant α in each orbit $[\alpha]$, that is, if one chooses a fixed section $\sigma : O \to S^{4\ell+1}$, $[\alpha] \to \sigma_{[\alpha]}$ with $[\sigma_{[\alpha]}] = [\alpha]$, every $\phi \in \mathbf{C}^{2\ell+1}$ can be uniquely written as

$$\phi = \rho\, U(R)\sigma_{[\alpha]}\ , \quad \rho = \|\phi\| \geq 0, \ \ \alpha = \tfrac{\phi}{\|\phi\|}, \quad [\alpha] \in O \text{ and } R \in SO(3)/I_{[\alpha]}$$

where $I_{[\alpha]} = I^{\sigma}_{[\alpha]} = \{S \in SO(3) \mid U(S)\sigma_{[\alpha]} = \sigma_{[\alpha]}\}$ is the isotropy subgroup of $\sigma_{[\alpha]}$. Let

$$\int_{\mathbf{R}^{4\ell+2}} \Pi_m\, du_m dv_m\, f(\phi) = \int_{\mathbf{R}^+} D\rho \int_O D[\alpha] \int_{[\alpha]} DR\, f\big(\rho\, U(R)\sigma_{[\alpha]}\big)$$

be the integral in (III.51) over $\mathbf{R}^{4\ell+2}$ in the new coordinates. That is, for example, $D\rho = \rho^{4\ell+1}d\rho$. In the new coordinates

$$|\Phi_{\mathbf{p}}|^2 = \lambda_\ell \rho^2 \Big|\sum_m \Big(\overline{U(R)\sigma_{[\alpha]}}\Big)_m Y_{\ell m}(\mathbf{p}')\Big|^2 = \lambda_\ell \rho^2 \Big|\sum_m \bar{\sigma}_{[\alpha],m} Y_{\ell m}\left(R^{-1}\mathbf{p}'\right)\Big|^2$$

such that

$$\tfrac{-a_{-p}}{a_p a_{-p} + |\Phi_{\mathbf{p}}|^2} = \tfrac{-a_{-p}}{a_p a_{-p} + \lambda_\ell \rho^2 |\sum_m \bar{\sigma}_{[\alpha],m} Y_{\ell m}(R^{-1}\mathbf{p}')|^2} \equiv f(\rho, [\alpha], R^{-1}\mathbf{p})$$

Since $V(\phi) = V(\rho, [\alpha])$ is independent of R, one obtains

$$\begin{aligned}
\tfrac{1}{\kappa}\langle \bar{\psi}_{p\sigma}\psi_{p\sigma}\rangle &= \frac{\int \tfrac{-a_{-p}}{a_p a_{-p} + |\Phi_{\mathbf{p}}|^2}\, e^{-\kappa V(\phi)} \prod\limits_{m=-\ell}^{\ell} du_m dv_m}{\int e^{-\kappa V(\phi)} \prod\limits_{m=-\ell}^{\ell} du_m dv_m} \\
&= \frac{\int_{\mathbf{R}^+} D\rho \int_O D[\alpha] \int_{[\alpha]} DR\ f(\rho, [\alpha], R^{-1}\mathbf{p})\ e^{-\kappa V(\rho,[\alpha])}}{\int_{\mathbf{R}^+} D\rho \int_O D[\alpha] \int_{[\alpha]} DR\ e^{-\kappa V(\rho,[\alpha])}} \\
&= \frac{\int_{\mathbf{R}^+} D\rho \int_O D[\alpha]\ \mathrm{vol}([\alpha])\ \frac{\int_{[\alpha]} DR f(\rho,[\alpha],R^{-1}\mathbf{p})}{\int_{[\alpha]} DR}\ e^{-\kappa V(\rho,[\alpha])}}{\int_{\mathbf{R}^+} D\rho \int_O D[\alpha]\ \mathrm{vol}([\alpha])\ e^{-\kappa V(\rho,[\alpha])}}
\end{aligned} \tag{III.53}$$

It is plausible to assume that at the global minimum of $V(\rho, [\alpha])$ ρ is uniquely determined, say ρ_0. Let $O_{\min} \subset O$ be the set of all orbits at which $V(\rho_0, [\alpha])$ takes its global minimum. Then in the infinite volume limit (III.53) becomes

$$\lim_{\kappa\to\infty} \tfrac{1}{\kappa}\langle \bar{\psi}_{p\sigma}\psi_{p\sigma}\rangle = \frac{\int_{O_{\min}} D[\alpha]\ \mathrm{vol}([\alpha])\ \frac{\int_{[\alpha]} DR\, f(\rho,[\alpha],R^{-1}\mathbf{p})}{\int_{[\alpha]} DR}}{\int_{O_{\min}} D[\alpha]\ \mathrm{vol}([\alpha])} \tag{III.54}$$

Consider the quotient of integrals in the numerator of (III.54). Since

$$\begin{aligned}
f(\rho, [\alpha], R^{-1}\mathbf{p}) &= f\Big(\rho^2 \big|\sum_m \bar{\sigma}_{[\alpha],m} Y_{\ell m}\left(R^{-1}\mathbf{p}'\right)\big|^2\Big) \\
&= f\Big(\rho^2 \big|\sum_m \big(\overline{U(S)\sigma_{[\alpha]}}\big)_m Y_{\ell m}\left(R^{-1}\mathbf{p}'\right)\big|^2\Big) \\
&= f\Big(\rho^2 \big|\sum_m \bar{\sigma}_{[\alpha],m} Y_{\ell m}\left((RS)^{-1}\mathbf{p}'\right)\big|^2\Big) = f(\rho, [\alpha], (RS)^{-1}\mathbf{p})
\end{aligned}$$

for all $S \in I_{[\alpha]}$, one has, since $[\alpha] \simeq SO(3)/I_{[\alpha]}$

$$\begin{aligned}
\frac{\int_{[\alpha]} DR\, f(\rho,[\alpha],R^{-1}\mathbf{p})}{\int_{[\alpha]} DR} &= \frac{\int_{SO(3)/I_{[\alpha]}} DR\, f(\rho,[\alpha],R^{-1}\mathbf{p}) \int_{I_{[\alpha]}} DS}{\int_{SO(3)/I_{[\alpha]}} DR \int_{I_{[\alpha]}} DS} \\
&= \frac{\int_{SO(3)/I_{[\alpha]}} DR \int_{I_{[\alpha]}} DS\, f(\rho,[\alpha],(RS)^{-1}\mathbf{p})}{\int_{SO(3)/I_{[\alpha]}} \int_{I_{[\alpha]}} DR\, DS} \\
&= \frac{\int_{SO(3)} DR\, f(\rho,[\alpha],R^{-1}\mathbf{p})}{\int_{SO(3)} DR} \\
&= \frac{\int_{S^2} d\Omega(\mathbf{x}) \int_{SO(3)_{\mathbf{x}\to\mathbf{p}}} DR\, f(\rho,[\alpha],R^{-1}\mathbf{p})}{\int_{S^2} d\Omega(\mathbf{x}) \int_{SO(3)_{\mathbf{x}\to\mathbf{p}}} DR} \\
&= \frac{\int_{S^2} d\Omega(\mathbf{x})\, f(\rho,[\alpha],\mathbf{x}) \int_{SO(3)_{\mathbf{x}\to\mathbf{p}}} DR}{\int_{S^2} d\Omega(\mathbf{x}) \int_{SO(3)_{\mathbf{x}\to\mathbf{p}}} DR} \qquad \text{(III.55)}
\end{aligned}$$

where $SO(3)_{\mathbf{x}\to\mathbf{p}} = \{R \in SO(3) \mid R\mathbf{x} = \mathbf{p}\}$. If one assumes that DR has the usual invariance properties of the Haar measure, then $\int_{SO(3)_{\mathbf{x}\to\mathbf{p}}} DR$ does not depend on $\mathbf{x}$ such that it cancells out in (III.55). Then (III.54) gives

$$\lim_{\kappa\to\infty} \tfrac{1}{\kappa}\langle \bar{\psi}_{p\sigma}\psi_{p\sigma}\rangle = \frac{\int_{O_{\min}} D[\alpha]\ \mathrm{vol}([\alpha])\ \frac{\int_{S^2} d\Omega(\mathbf{x})\, f(\rho,[\alpha],\mathbf{x})}{\int_{S^2} d\Omega(\mathbf{x})}}{\int_{O_{\min}} D[\alpha]\ \mathrm{vol}([\alpha])} \qquad \text{(III.56)}$$

Now, since the effective potential, which is constant on $O_{\min}$, may be written as

$$V(\rho,[\alpha]) = \int_{S^2} \tfrac{d\Omega(\mathbf{x})}{4\pi}\, G\Big(\rho^2 \big|\Sigma_m\, \bar{\sigma}_{[\alpha],m} Y_{\ell m}(\mathbf{x})\big|^2\Big)$$

with $G(X) = \rho^2 - \int \frac{dk\,k^2}{2\pi^2}\, \log\left[\frac{\cosh(\frac{\beta}{2}\sqrt{e_k^2+\lambda_\ell X})}{\cosh\frac{\beta}{2}e_k}\right]$, it is plausible to assume that also

$$\frac{\int_{S^2} d\Omega(\mathbf{x})\, f(\rho,[\alpha],\mathbf{x})}{\int_{S^2} d\Omega(\mathbf{x})} = \int_{S^2} \frac{d\Omega(\mathbf{x})}{4\pi}\, \frac{ip_0+e_{\mathbf{p}}}{p_0^2+e_{\mathbf{p}}^2+\lambda_\ell\rho^2|\,\Sigma_m\, \bar{\sigma}_{[\alpha],m} Y_{\ell m}(x)|^2}$$

is constant on $O_{\min}$. In that case also the integrals over $O_{\min}$ in (III.56) cancel out and the theorem is proven •

IV. Solution with Delta Function Interaction and Its Perturbation Theory

IV.1 Solution with Delta Function

We consider the model

$$Z(\beta,L,\{s_{k,\sigma}\}) = \int e^{-\mathcal{U}(\psi,\bar\psi)+\frac{1}{\kappa}\sum_{k,\sigma} s_{k,\sigma}\bar\psi_{k,\sigma}\psi_{k,\sigma}}\, d\mu_C(\psi,\bar\psi) \tag{IV.1}$$

where

$$\begin{aligned}\mathcal{U}(\psi,\bar\psi) &= \tfrac{\lambda}{\kappa^3}\sum_{\sigma,\tau\in\{\uparrow,\downarrow\}}\sum_{k,p,q}\left[\delta_{k,p}+\delta_{k,q-p}+\delta_{q,0}\right]\bar\psi_{k\sigma}\bar\psi_{q-k\tau}\psi_{p\sigma}\psi_{q-p\tau}\\ &= \tfrac{\lambda}{\kappa^3}\sum_{k,p,q}\left[\delta_{k,p}+\delta_{k,q-p}+\delta_{q,0}\right]\bar\psi_{k\uparrow}\bar\psi_{q-k\downarrow}\psi_{p\uparrow}\psi_{q-p,\downarrow}\end{aligned} \tag{IV.2}$$

It is useful to explicitly cancel the $\bar\psi_\uparrow\bar\psi_\uparrow\psi_\uparrow\psi_\uparrow$ and $\bar\psi_\downarrow\bar\psi_\downarrow\psi_\downarrow\psi_\downarrow$ terms before directly applying Theorem II.1 because then the integral representation becomes 4 dimensional instead of 5 dimensional. Both representations are of course equivalent. One obtains

$$\langle\bar\psi_{p\sigma}\psi_{p'\sigma}\rangle = \beta L^d\delta_{p,p'}\,\frac{\int \frac{\frac{\partial}{\partial s_{p\sigma}}\mathrm{Pf}S_p}{\mathrm{Pf}S_p}\, e^{-\kappa V_\beta(z,x,\rho)}dv\,dw\,xdx\,\rho d\rho}{\int e^{-\kappa V_\beta(z,x,\rho)}dv\,dw\,xdx\,\rho d\rho} \tag{IV.3}$$

where $z=v+iw$,

$$\begin{aligned}\mathrm{Pf}S_k =& (A_{k\uparrow}A_{-k\downarrow}+\lambda\rho^2)(A_{-k\uparrow}A_{k\downarrow}+\lambda\rho^2)+(A_{k\uparrow}A_{k\downarrow}+A_{-k\uparrow}A_{-k\downarrow})\lambda x^2\\ &+2\lambda^2x^2\rho^2+\lambda^2x^4\end{aligned} \tag{IV.4}$$

$$A_{k\uparrow}=a_k-s_{k\uparrow}-\sqrt{\lambda}\,z,\qquad A_{k\downarrow}=a_k-s_{k\downarrow}-\sqrt{\lambda}\,\bar z$$

as before $a_k=ik_0-e_{\mathbf{k}}$ and the effective potential is given by

$$V_\beta(z,x,\rho)=|z|^2+x^2+\rho^2-\tfrac{1}{\beta L^d}\sum_{\mathbf{k}\in M_\omega}\log\prod_{k_0>0}\tfrac{\mathrm{Pf}S_k}{a_k^2a_{-k}^2} \tag{IV.5}$$

The product over k_0 is computed in Lemma A1 in the appendix. One finds

$$V_\beta(v,w,x,\rho)= \tag{IV.6}$$
$$v^2+w^2+x^2+\rho^2-\tfrac{1}{L^d}\sum_{\mathbf{k}\in M_\omega}\tfrac{1}{\beta}\log\left[\frac{\cosh^2\left(\frac{\beta}{2}\sqrt{(e_{\mathbf{k}}+\sqrt{\lambda}v)^2+\lambda\rho^2}\right)-\sin^2\left(\frac{\beta}{2}\sqrt{\lambda(w^2+x^2)}\right)}{\cosh^2\frac{\beta}{2}e_{\mathbf{k}}}\,e^{-\beta\sqrt{\lambda}v}\right]$$

The z variable sums up forward contributions, $x=|\xi|$ sums up exchange contributions and $\rho=|\phi|$ collects the BCS contributions. Pure BCS is given by $z=x=0$ and (IV.6) coincides with (I.47) or (III.10) (for $r=0$).

To compute the infinite volume limit of (IV.3) one has to find the global minimum of the effective potential (IV.6).

Consider first the case of an attractive coupling $\lambda = g^2 > 0$. To make (IV.6) small, the numerator in the logarithm has to be large. Hence for λ positive $w^2 + x^2$ has to be zero. This gives

$$\frac{\frac{\partial}{\partial s_{p\uparrow}}\mathrm{Pf}S_p}{\mathrm{Pf}S_p} = \frac{a_{-p}-gv}{[a_p-gv][a_{-p}-gv]+g^2\rho^2} \tag{IV.7}$$

As in section III.1 ρ is positive for sufficiently small $T = \frac{1}{\beta}$ and v may or may not be nonzero to renormalize $e_{\mathbf{p}}$. Thus for attractive coupling the two point function becomes

$$\lim_{L\to\infty} \tfrac{1}{\kappa}\langle\bar\psi_{p\uparrow}\psi_{p\uparrow}\rangle = -\frac{ip_0+e_{\mathbf{p}g}}{p_0^2+e_{\mathbf{p}g}^2+|\Delta|^2} \tag{IV.8}$$

if $e_{\mathbf{p}g} = e_{\mathbf{p}} + gv_0$, $|\Delta|^2 = \lambda\rho_0^2$ and v_0, ρ_0 are the (λ dependent) values where V takes its global minimum.

Now consider a repulsive coupling $\lambda = -g^2 < 0$. In that case the numerator in the logarithm in (IV.6) reads $\cosh^2(\frac{\beta}{2}\sqrt{(e_{\mathbf{k}}+igv)^2 - g^2\rho^2}\,) + \sinh^2(\frac{\beta}{2}g\sqrt{w^2+x^2}\,)$ and the value $\rho = 0$ is favourable. Furthermore $v = 0$ seems favourable. In that case

$$\frac{\frac{\partial}{\partial s_{p\uparrow}}\mathrm{Pf}S_p}{\mathrm{Pf}S_p} = \frac{a_p-gw}{a_p^2-g^2(w^2+x^2)} \tag{IV.9}$$

and the two point function becomes, if $\delta_g^2 = g^2(w^2+x^2) \geq 0$

$$\lim_{L\to\infty} \tfrac{1}{\kappa}\langle\bar\psi_{p\uparrow}\psi_{p\uparrow}\rangle = \frac{a_p}{a_p^2-\delta_g^2} = \frac{1}{2}\left[\frac{1}{ip_0-e_{\mathbf{p}}-\delta_g} + \frac{1}{ip_0-e_{\mathbf{p}}+\delta_g}\right] \tag{IV.10}$$

since the effective potential is even in w and therefore the w term in (IV.9) cancels. At zero temperature, this results in a momentum distribution

$$\begin{aligned} n_{\mathbf{k}} &= \lim_{\substack{L\to\infty\\ \beta\to\infty}} \tfrac{1}{L^d}\langle a_{\mathbf{k}}^+ a_{\mathbf{k}}\rangle_{\beta,L} = \lim_{\substack{\epsilon\to 0\\ \epsilon<0}} \int_{\mathbf{R}} \frac{dk_0}{2\pi}\frac{1}{2}\left[\frac{1}{e^{ik_0\epsilon}(ik_0-e_{\mathbf{k}})-\delta_g} + \frac{1}{e^{ik_0\epsilon}(ik_0-e_{\mathbf{k}})+\delta_g}\right] \\ &= \tfrac{1}{2}\Big[\chi(e_{\mathbf{k}}-\delta_g<0) + \chi(e_{\mathbf{k}}+\delta_g<0)\Big] = \begin{cases} 1 & \text{if } e_{\mathbf{k}} < -\delta_g \\ \frac{1}{2} & \text{if } -\delta_g < e_{\mathbf{k}} < \delta_g \\ 0 & \text{if } \delta_g < e_{\mathbf{k}} \end{cases} \end{aligned} \tag{IV.11}$$

Thus, if δ_g is nonzero, the exchange and forward contributions lead to a splitting of the Fermi surface. A nonzero δ_g can be achieved by making the coupling g^2 sufficiently big: Let $K = \frac{|S^{d-1}|}{(2\pi)^d} m k_F^{d-2} \sim k_F^d/\mu$ and let

$$V_\beta(y) = y^2 - \int_{M_\omega} \frac{d^d\mathbf{k}}{(2\pi)^d}\,\frac{1}{\beta}\log\left[\frac{\cosh^2(\frac{\beta}{2}e_{\mathbf{k}}) + \sinh^2(\frac{\beta}{2}gy)}{\cosh^2\frac{\beta}{2}e_{\mathbf{k}}}\right] \tag{IV.12}$$

Then

$$\lim_{\beta\to\infty} V_\beta(y) = (1-Kg^2)y^2 + K[gy-\omega]_+^2 \tag{IV.13}$$

where $[x]_+ = x$ for positive x and 0 otherwise. The global minimum of V_∞ is at $y_{\min}$ where

$$y_{\min} = \begin{cases} 0 & \text{if } g^2 < 1/K \\ K\omega g & \text{if } g^2 > 1/K \end{cases} \tag{IV.14}$$

IV.2 Perturbation Theory

In this section we consider the perturbation theory of the model

$$Z_{\text{ex/BCS}}(\beta, L) = \int e^{-\mathcal{V}_{\text{ex/BCS}}(\psi,\bar\psi)} d\mu_C(\psi,\bar\psi) \tag{IV.15}$$

For BCS, one obtains a power series in $C(k)C(-k)$ and for $\mathcal{V}_{\text{ex}}$ one gets a power series in $C(k)^2$.

We will find that the linked cluster theorem, $\log Z$ is given by the sum of all connected diagrams, cannot be applied to (IV.15). More specifically, whereas the series for $Z(\lambda)$, the sum of all diagrams, converges for sufficiently small λ (for finite β and L) with $Z(0) = 1$ which implies that also $\log Z$ is analytic for λ sufficiently small, the sum of all connected diagrams has radius of convergence zero (for finite β and L).

We start with BCS. First we show how the integral representation (III.12) (for $r = 0$)

$$Z_{\text{BCS}} = 2\kappa \int e^{-\kappa\left(\rho^2 - \frac{1}{L^d}\sum_{\mathbf{k}} \log\left[\frac{\cosh(\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\lambda\rho^2})}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\right]\right)} \rho\, d\rho \tag{IV.16}$$

is obtained by direct summation of the diagrams without making a Hubbard Stratonovich transformation. That is, without using the identity

$$e^{-\frac{\lambda}{\kappa^3}\sum_{k,p}\psi_{k\uparrow}\psi_{-k\downarrow}\bar\psi_{p,\uparrow}\psi_{-p\downarrow}} = \int e^{i\phi\, g\frac{1}{\kappa}\sum_k \psi_{k\uparrow}\psi_{-k\downarrow} + i\bar\phi\, g\frac{1}{\kappa}\sum_k \bar\psi_{k\uparrow}\bar\psi_{-k\downarrow}} \frac{\kappa}{\pi}\, e^{-\kappa|\phi|^2} dudv$$

One has

$$\begin{aligned} Z_{\text{BCS}} &= \int e^{-\frac{\lambda}{\kappa^3}\sum_{k,p}\psi_{k\uparrow}\psi_{-k\downarrow}\bar\psi_{p\uparrow}\bar\psi_{-p\downarrow}} d\mu_C(\psi,\bar\psi) \qquad \text{(IV.17)} \\ &= \sum_{n=0}^\infty \frac{\left(\frac{\lambda}{\kappa^3}\right)^n}{n!} \sum_{\substack{k_1,\cdots,k_n \\ p_1,\cdots,p_n}} \det\left[\kappa\delta_{k_i,p_j}C(k_i)\right]_{1\le i,j\le n} \det\left[\kappa\delta_{k_i,p_j}C(-k_i)\right]_{1\le i,j\le n} \\ &= \sum_{n=0}^\infty \left(\tfrac{\lambda}{\kappa}\right)^n \sum_{\pi\in S_n} \epsilon_\pi \sum_{k_1,\cdots,k_n} C(k_1)C(-k_1)\cdots C(k_n)C(-k_n)\,\delta_{k_1,k_{\pi 1}}\cdots\delta_{k_n,k_{\pi n}} \end{aligned}$$

This is the expansion into Feynman diagrams. It can be summed up by collecting the fermion loops:

Say that the permutation π is of type $t(\pi) = 1^{b_1} \cdots n^{b_n}$ if the decomposition into disjoint cycles contains b_r r-cycles for $1 \le r \le n$. Necessarily one has $1b_1 + \cdots + nb_n = n$. The number of permutations which have b_r r-cycles for $1 \le r \le n$ is

$$\frac{n!}{b_1! \cdots b_n! \, 1^{b_1} \cdots n^{b_n}}$$

The sign of such a permutation is given by

$$\epsilon_\pi = (-1)^{(1-1)b_1 + (2-1)b_2 + \cdots + (n-1)b_n} = (-1)^{n - \sum_{r=1}^n b_r}$$

Therefore one obtains

$$\begin{aligned} Z = \sum_{n=0}^{\infty} \left(\tfrac{\lambda}{\kappa}\right)^n \sum_{\substack{b_1,\cdots,b_n=0 \\ 1b_1+\cdots+nb_n=n}}^{n} & \frac{n!}{b_1! \cdots b_n! \, 1^{b_1} \cdots n^{b_n}} (-1)^{n-\sum_{r=1}^n b_r} \times \\ & \Big\{ \sum_k [C(k)C(-k)]^1 \Big\}^{b_1} \cdots \Big\{ \sum_k [C(k)C(-k)]^n \Big\}^{b_n} \\ = \sum_{n=0}^{\infty} \sum_{\substack{b_1,\cdots,b_n=0 \\ 1b_1+\cdots+nb_n=n}}^{n} & n! \prod_{r=1}^{n} \frac{1}{b_r!} \Big\{ -\tfrac{1}{r} \sum_k \left[-\tfrac{\lambda}{\kappa} C(k)C(-k) \right]^r \Big\}^{b_r} \end{aligned} \tag{IV.18}$$

The only factor which prevents us from an explicit summation of the above series is the $n!$. Therefore we substitute

$$n! = \int_0^\infty e^{-x} x^n dx$$

and obtain

$$\begin{aligned} Z &= \int_0^\infty e^{-x} \sum_{n=0}^{\infty} \sum_{\substack{b_1,\cdots,b_n=0 \\ 1b_1+\cdots+nb_n=n}}^{n} \prod_{r=1}^{n} \frac{1}{b_r!} \Big\{ -\tfrac{1}{r} \sum_k \left[-\tfrac{\lambda}{\kappa} x \, C(k)C(-k) \right]^r \Big\}^{b_r} dx \\ &= \int_0^\infty e^{-x} \prod_{r=1}^{\infty} \sum_{b_r=0}^{\infty} \frac{1}{b_r!} \Big\{ -\tfrac{1}{r} \sum_k \left[-\tfrac{\lambda}{\kappa} x \, C(k)C(-k) \right]^r \Big\}^{b_r} dx \\ &= \int_0^\infty e^{-x} e^{-\sum_k \sum_{r=1}^\infty \frac{1}{r} \left[-\frac{\lambda}{\kappa} x \, C(k)C(-k) \right]^r} dx \\ &= \int_0^\infty e^{-x} e^{\sum_k \log\left[1 + \frac{\lambda x}{\kappa} \, C(k)C(-k) \right]} dx \\ &= 2\kappa \int_0^\infty e^{-\kappa\left(\rho^2 - \frac{1}{\kappa} \sum_k \log\left[1 + \frac{\lambda \rho^2}{k_0^2 + e_{\mathbf{k}}^2} \right] \right)} \rho \, d\rho \\ &= 2\kappa \int_0^\infty e^{-\kappa\left(\rho^2 - \frac{1}{L^d} \sum_{\mathbf{k}} \frac{1}{\beta} \log\left[\frac{\cosh(\frac{\beta}{2} \sqrt{e_{\mathbf{k}}^2 + \lambda\rho^2})}{\cosh \frac{\beta}{2} e_{\mathbf{k}}} \right]^2 \right)} \rho \, d\rho \end{aligned} \tag{IV.19}$$

which coincides with (IV.16). In the last line we used (III.25) again. The case of an exchange interaction is treated in the same way:

$$
\begin{aligned}
Z_{\mathrm{ex}} &= \int e^{-\frac{\lambda}{\kappa^3}\sum_{k,p}\psi_{k\uparrow}\psi_{p\downarrow}\bar\psi_{p\uparrow}\bar\psi_{k\downarrow}}\,d\mu_C(\psi,\bar\psi) \\
&= \sum_{n=0}^{\infty}\left(\tfrac{\lambda}{\kappa}\right)^n \sum_{\pi\in S_n}\epsilon_\pi \sum_{k_1,\cdots,k_n} C(k_1)^2\cdots C(k_n)^2\,\delta_{k_1,k_{\pi 1}}\cdots\delta_{k_n,k_{\pi n}} \\
&= \sum_{n=0}^{\infty}\sum_{\substack{b_1,\cdots,b_n=0\\ 1b_1+\cdots+nb_n=n}}^{n} n!\prod_{r=1}^{n}\frac{1}{b_r!}\Big\{-\tfrac{1}{r}\sum_k\left[-\tfrac{\lambda}{\kappa}C(k)^2\right]^r\Big\}^{b_r} && \text{(IV.20)} \\
&= \int_0^\infty e^{-x} e^{-\sum_k\sum_{r=1}^\infty\frac{1}{r}\left[-\frac{\lambda}{\kappa}x\,C(k)^2\right]^r}\,dx \\
&= 2\kappa\int_0^\infty e^{-\kappa\left(y^2-\frac{1}{L^d}\sum_{\mathbf{k}}\frac{1}{\beta}\log\left[\frac{\cosh^2(\frac{\beta}{2}e_{\mathbf{k}})-\sin^2(\frac{\beta}{2}\sqrt{\lambda}\,y)}{\cosh^2\frac{\beta}{2}e_{\mathbf{k}}}\right]^2\right)}\,y\,dy && \text{(IV.21)}
\end{aligned}
$$

where in the last line Lemma A1 in the appendix has been used to compute the k_0 product. Observe that $\lim_{\beta\to\infty}\frac{1}{\kappa}\sum_k C(k)^{2r}=0$ for all $r\ge 1$ since $\int\frac{dk_0}{2\pi}\frac{1}{(ik_0-e)^j}=0$ for all $j\ge 2$, but for large repulsive coupling $\lim_{\beta\to\infty}\frac{1}{\beta L^d}\log Z_{\mathrm{ex}}=-V_\infty(y_{\min})>0$ by (IV.13,14).

We now consider the linked cluster theorem. It states that, if the partition function

$$Z(\lambda)=\sum_{n=0}^{\infty}\lambda^n\sum_{G\in\Gamma_n}\mathrm{val}(G) \qquad \text{(IV.22)}$$

is given by a sum of diagrams, Γ_n being the set of all n'th order diagrams, then the logarithm

$$\log Z(\lambda)=\sum_{n=0}^{\infty}\lambda^n\sum_{G\in\Gamma_n^c}\mathrm{val}(G) \qquad \text{(IV.23)}$$

is given by the sum of all connected diagrams. This theorem is easily illustrated for a quadratic perturbation. Namely, if

$$
\begin{aligned}
Z &= \int e^{-\frac{\lambda}{\kappa}\sum_{k\sigma}\bar\psi_{k\sigma}\psi_{k\sigma}}\,d\mu_C(\psi,\bar\psi) \\
&= \sum_{n=0}^{\infty}\frac{(-\lambda)^n}{n!}\sum_{\substack{k_1\cdots k_n\\ \sigma_1\cdots\sigma_n}}\det\left[\delta_{\sigma_i,\sigma_j}\delta_{k_i,k_j}C(k_i)\right]_{1\le i,j\le n} \\
&= \sum_{n=0}^{\infty}\sum_{\substack{b_1,\cdots,b_n=0\\ 1b_1+\cdots+nb_n=n}}^{n}\prod_{r=1}^{n}\frac{1}{b_r!}\Big\{-\tfrac{1}{r}\sum_k\left[-\lambda C(k)\right]^r\Big\}^{b_r} && \text{(IV.24)} \\
&= e^{\sum_k\log[1+\lambda C(k)]} && \text{(IV.25)}
\end{aligned}
$$

then the sum of all connected diagrams is obtained by summing all the terms with $b_1 = \cdots = b_{n-1} = 0$ and $b_n = 1$ in (IV.24). That is, one gets

$$\sum_{n=1}^{\infty} \frac{1}{1!} \Big\{ -\tfrac{1}{n} \sum_k \left[-\lambda C(k)\right]^n \Big\}^1 = \sum_k \log\left[1 + \lambda C(k)\right] \tag{IV.26}$$

which coincides with $\log Z$. Now consider $Z_{\rm BCS}$. The sum of all connected diagrams is given by the sum of all the terms with $b_1 = \cdots = b_{n-1} = 0$ and $b_n = 1$ in (IV.18). That is, one obtains

$$\sum_{n=0}^{\infty} (-1)^{n+1} \left(\tfrac{\lambda}{\kappa}\right)^n n!\, \tfrac{1}{n} \sum_k \left[C(k)C(-k)\right]^n \tag{IV.27}$$

which has radius of convergence zero, at finite temperature and finite volume. However, $Z_{\rm BCS}(\lambda)$ has positive radius of convergence and $Z_{\rm BCS}(0) = 1$ which means that also $\log Z_{\rm BCS}(\lambda)$ is analytic for sufficiently small (volume and temperature dependent) λ. That is, in this case the linked cluster theorem does not apply since the right hand side of (IV.23) is infinite.

We remark that we think that this is an artefact of the specific model at hand for the following reason. Suppose for the moment that the k sums in (IV.17,18) are finite with N different values of k. Then $Z_{\rm BCS}(\lambda)$ is a polynomial in λ of degree (at most) N. In particular, the coefficients

$$\sum_{\substack{b_1,\cdots,b_n=0 \\ 1b_1+\cdots+nb_n=n}}^{n} \; \prod_{r=1}^{n} \frac{1}{b_r!} \Big\{ -\tfrac{1}{r} \sum_k \left[C(k)C(-k)\right]^r \Big\}^{b_r} \;=\; 0 \quad \text{if} \;\; n > N\,. \tag{IV.28}$$

That is, there are strong cancellations between fermion loops of different orders. However, for $\mathcal{V}_{\rm int} \approx \mathcal{V}_{\rm BCS}$ (or forward or exchange), an n'th order connected diagram is necessarily a single n'th order fermion loop such that there are no cancellations at all. In a more realistic model a connected diagram contains fermion loops of different orders and cancellations are present. In fact, a careful diagrammatic analysis [FT,FKLT,FST] shows that the renormalized sum of all connected diagrams of the two dimensional electron system with anisotropic dispersion relation $e_{\mathbf{k}}$ (such that Cooper pairs are suppressed) has positive radius of convergence (which is, in particular, independent of volume and temperature).

Appendix

Lemma A.1: a) *Let* $a_k = ik_0 - e_{\mathbf{k}}$, $e_{-\mathbf{k}} = e_{\mathbf{k}}$ *and let* b, c, d *be some complex numbers. Then*

$$\prod_{k_0 \in \frac{\pi}{\beta}(2\mathbf{Z}+1)} \tfrac{a_k a_{-k} + b a_k + c a_{-k} + d}{a_k a_{-k}} \;:=\; \lim_{\substack{\epsilon \to 0 \\ \epsilon < 0}} \prod_{k_0 \in \frac{\pi}{\beta}(2\mathbf{Z}+1)} \tfrac{a_k a_{-k} + b e^{i\epsilon k_0} a_k + c e^{-i\epsilon k_0} a_{-k} + d}{a_k a_{-k}}$$

$$= \tfrac{\cosh\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\omega_+}}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\;\tfrac{\cosh\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\omega_-}}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\;e^{\frac{\beta}{2}(b+c)} \tag{A.1}$$

where $\omega_\pm = \frac{p\pm\sqrt{p^2-4q}}{2}$ *with* $p = (b+c-e_{\mathbf{k}})^2 - e_k^2 - 4bc + 2d$ *and* $q = 4bce_{\mathbf{k}}^2 - 2(b+c)e_{\mathbf{k}}d + d^2$*. In particular, for* $b = c$ *and* $d = c^2$

$$\prod_{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)} \left(1+\tfrac{c}{a_k}\right) := \lim_{\substack{\epsilon\to 0\\ \epsilon<0}} \prod_{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)} \left(1+e^{-i\epsilon k_0}\tfrac{c}{a_k}\right) = \tfrac{1+e^{-\beta(e_{\mathbf{k}}-c)}}{1+e^{-\beta e_{\mathbf{k}}}} \tag{A.2}$$

b) *Let* $A_{k\uparrow} = a_k - \sqrt{\lambda}\,z$*,* $A_{k\downarrow} = a_k - \sqrt{\lambda}\,\bar{z}$*, and let*

$$\begin{aligned}\mathrm{Pf}S_k = \mathrm{Pf}S_k(a_k,a_{-k}) &= (A_{k\uparrow}A_{-k\downarrow}+\lambda\rho^2)(A_{k\downarrow}A_{-k\uparrow}+\lambda\rho^2)\\ &\quad + (A_{k\uparrow}A_{k\downarrow}+A_{-k\uparrow}A_{-k\downarrow}+2\lambda\rho^2)\lambda x^2+\lambda^2x^4\end{aligned} \tag{A.3}$$

Then

$$\begin{aligned}\prod_{\substack{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)\\ k_0>0}} \tfrac{\mathrm{Pf}S_k(a_k,a_{-k})}{a_k^2a_{-k}^2} &:= \lim_{\substack{\epsilon\to 0\\ \epsilon<0}} \prod_{\substack{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)\\ k_0>0}} \tfrac{\mathrm{Pf}S_k(e^{i\epsilon k_0}a_k,e^{-i\epsilon k_0}a_{-k})}{a_k^2a_{-k}^2}\\ &= \tfrac{\cosh^2\left(\frac{\beta}{2}\sqrt{(e_{\mathbf{k}}+\sqrt{\lambda}\,v)^2+\lambda\rho^2}\right)-\sin^2\left(\frac{\beta}{2}\sqrt{\lambda(w^2+x^2)}\right)}{\cosh^2\frac{\beta}{2}e_{\mathbf{k}}}\;e^{-\beta\sqrt{\lambda}v}\end{aligned} \tag{A.4}$$

Proof: We first explain why the definition of the product given in the lemma is the right one.

Consider the quadratic perturbation

$$H = H_0 + \lambda N = \tfrac{1}{L^d}\sum_{\mathbf{k}\sigma} e_{\mathbf{k}}\,a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma} + \tfrac{\lambda}{L^d}\sum_{\mathbf{k}\sigma} a^+_{\mathbf{k}\sigma}a_{\mathbf{k}\sigma}$$

One finds by explicit computation of the trace

$$Z(\lambda,\beta) = \tfrac{Tr\,e^{-\beta(H_0+\lambda N)}}{Tr\,e^{-\beta H_0}} = \prod_{\mathbf{k}\sigma} \tfrac{1+e^{-\beta(e_{\mathbf{k}}+\lambda)}}{1+e^{-\beta e_{\mathbf{k}}}}$$

On the other hand, by explicit summation of the perturbation series or equivalently, by performing the fermionic functional integral, one obtains

$$\begin{aligned}Z(\lambda,\beta) &= \int e^{\frac{\lambda}{\beta L^d}\sum_{k\sigma}\bar{\psi}_{k\sigma}\psi_{k\sigma}}d\mu_C\\ &= \prod_{k\sigma}\tfrac{\beta L^d}{ik_0-e_{\mathbf{k}}}\int e^{\frac{\lambda}{\beta L^d}\sum_{k\sigma}\bar{\psi}_{k\sigma}\psi_{k\sigma}}e^{-\frac{1}{\beta L^d}\sum_{k\sigma}(ik_0-e_{\mathbf{k}})\bar{\psi}_{k\sigma}\psi_{k\sigma}}\prod_{k\sigma}d\psi_{k\sigma}d\bar{\psi}_{k\sigma}\\ &= \prod_{k\sigma}\tfrac{ik_0-e_{\mathbf{k}}-\lambda}{ik_0-e_{\mathbf{k}}} = \prod_{\mathbf{k}\sigma}\prod_{k_0}\left(1-\tfrac{\lambda}{a_k}\right)\end{aligned} \tag{A.5}$$

In the perturbation expansion the product in the last equation results from

$$\prod_{k\sigma}\left(1-\tfrac{\lambda}{a_k}\right)=e^{\sum_{k\sigma}\log\left(1-\frac{\lambda}{a_k}\right)}=e^{-\sum_{r=1}^{\infty}\frac{\lambda^r}{r}\sum_{k\sigma}\frac{1}{{a_k}^r}}$$

The term for $r=1$ requires special attention since $\sum_{k_0}\frac{1}{ik_0-e_{\mathbf{k}}}$ is not absolutely convergent. This sum enters the perturbation series as the propagator

$$C(x_0-x_0',\mathbf{x}-\mathbf{x}')=\tfrac{Tr\,e^{-\beta H_0}T\psi(x_0\mathbf{x})\psi^+(x_0'\mathbf{x}')}{Tr\,e^{-\beta H_0}}$$

evaluated at $x_0-x_0'=0$. Here $T\psi(x_0\mathbf{x})\psi^+(x_0'\mathbf{x}')$ is defined to be $\psi(x_0\mathbf{x})\psi^+(x_0'\mathbf{x}')$ if $x_0\geq x_0'$ and $-\psi^+(x_0'\mathbf{x}')\psi(x_0\mathbf{x})$ if $x_0<x_0'$. By explicit computation one finds that for $x_0\neq x_0'$

$$\begin{aligned}C(x_0-x_0',\mathbf{x}-\mathbf{x}')&=\tfrac{1}{L^d}\sum_{\mathbf{k}}e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')}e^{-e_{\mathbf{k}}(x_0-x_0')}\left[\theta_\beta(-e_{\mathbf{k}})\theta(x_0'-x_0)-\theta_\beta(e_{\mathbf{k}})\theta(x_0-x_0')\right]\\&=\tfrac{1}{\beta L^d}\sum_{\mathbf{k},k_0}e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')-ik_0(x_0-x_0')}\tfrac{1}{ik_0-e_{\mathbf{k}}}\end{aligned}$$

and for $x_0=x_0'$

$$C(0,\mathbf{x}-\mathbf{x}')=\tfrac{1}{L^d}\sum_{\mathbf{k}}e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')}\theta_\beta(-e_{\mathbf{k}})=\lim_{\substack{\epsilon\to0\\ \epsilon<0}}\tfrac{1}{\beta L^d}\sum_{\mathbf{k},k_0}e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')-ik_0\epsilon}\tfrac{1}{ik_0-e_{\mathbf{k}}}$$

Here θ is a step function being 1 for positive arguments and $\theta_\beta(-v)=\frac{e^{-\beta v}}{1+e^{-\beta v}}$ is an approximate step function. Therefore the k_0-sums have to be evaluated according to

$$\sum_{k_0}\tfrac{1}{ik_0-e_{\mathbf{k}}}=\sum_{k_0}^{\epsilon}\tfrac{1}{ik_0-e_{\mathbf{k}}}:=\lim_{\substack{\epsilon\to0\\ \epsilon<0}}e^{-ik_0\epsilon}\tfrac{1}{ik_0-e_{\mathbf{k}}}=\beta\theta_\beta(-e_{\mathbf{k}})=\beta\,\tfrac{e^{-\beta e_{\mathbf{k}}}}{1+e^{-\beta e_{\mathbf{k}}}}$$

We emphasize this point because the following computation leads to a wrong result

$$\sum_{k_0}\tfrac{1}{ik_0-e_{\mathbf{k}}}=\sum_{k_0}^{sym}\tfrac{1}{ik_0-e_{\mathbf{k}}}:=\tfrac{1}{2}\sum_{k_0}\left[\tfrac{1}{ik_0-e_{\mathbf{k}}}+\tfrac{1}{-ik_0-e_{\mathbf{k}}}\right]=\sum_{k_0}\tfrac{-e_{\mathbf{k}}}{k_0^2+e_{\mathbf{k}}^2}=-\tfrac{\beta}{2}\,\tfrac{1-e^{-\beta e_{\mathbf{k}}}}{1+e^{-\beta e_{\mathbf{k}}}}$$

Therefore the product in (A.5) is found to be

$$\begin{aligned}\prod_{k_0}\left(1-\tfrac{\lambda}{ik_0-e_{\mathbf{k}}}\right)&=\prod_{k_0}^{\epsilon}\left(1-\tfrac{\lambda}{ik_0-e_{\mathbf{k}}}\right)=e^{-\sum_{r=1}^{\infty}\frac{\lambda^r}{r}\sum_{k_0}^{\epsilon}\frac{1}{(ik_0-e_{\mathbf{k}})^r}}\\&=e^{-\sum_{r=1}^{\infty}\frac{\lambda^r}{r}\sum_{k_0}^{sym}\frac{1}{(ik_0-e_{\mathbf{k}})^r}}\;e^{\sum_{k_0}^{sym}\frac{\lambda}{ik_0-e_{\mathbf{k}}}-\sum_{k_0}^{\epsilon}\frac{\lambda}{ik_0-e_{\mathbf{k}}}}\\&=\prod_{k_0}^{sym}\left(1-\tfrac{\lambda}{ik_0-e_{\mathbf{k}}}\right)e^{-\frac{\beta}{2}\lambda}=\prod_{k_0>0}\tfrac{k_0^2+(e_{\mathbf{k}}+\lambda)^2}{k_0^2+e_{\mathbf{k}}^2}\;e^{-\frac{\beta}{2}\lambda}\\&\overset{(III.25)}{=}\tfrac{\cosh\frac{\beta}{2}(e_{\mathbf{k}}+\lambda)}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\;e^{-\frac{\beta}{2}\lambda}=\tfrac{1+e^{-\beta(e_{\mathbf{k}}+\lambda)}}{1+e^{-\beta e_{\mathbf{k}}}}\end{aligned}\tag{A.6}$$

which proves (A.2).

In the general case one may proceed similarly to obtain (using $a_k + a_{-k} = -2e_{\mathbf{k}}$)

$$\begin{aligned}
\prod_{k_0 \in \frac{\pi}{\beta}(2\mathbf{Z}+1)} \tfrac{a_k a_{-k} + b a_k + c a_{-k} + d}{a_k a_{-k}} &= \lim_{\substack{\epsilon \to 0 \\ \epsilon < 0}} \prod_{k_0} \left(1 + \tfrac{b e^{i\epsilon k_0} a_k + c e^{-i\epsilon k_0} a_{-k} + d}{a_k a_{-k}}\right) =: \prod_{k_0}^{\epsilon} \cdots \\
&= \prod_{k_0}^{sym} \left(1 + \tfrac{b a_k + c a_{-k} + d}{a_k a_{-k}}\right) e^{b \sum_{k_0}^{\epsilon} - \sum_{k_0}^{sym} \frac{1}{a_{-k}} + c \sum_{k_0}^{\epsilon} - \sum_{k_0}^{sym} \frac{1}{a_k}} \\
&= \prod_{k_0 > 0} \tfrac{(a_k a_{-k} + b a_k + c a_{-k} + d)(a_k a_{-k} + b a_{-k} + c a_k + d)}{(a_k a_{-k})^2} \; e^{\frac{\beta}{2}(b+c)} \\
&= \prod_{k_0 > 0} \tfrac{(a_k a_{-k})^2 + a_k a_{-k}(-2e_{\mathbf{k}}(b+c) + 2d + b^2 + c^2) + bc(a_k^2 + a_{-k}^2) - 2e_{\mathbf{k}} d(b+c) + d^2}{(a_k a_{-k})^2} \; e^{\frac{\beta}{2}(b+c)}
\end{aligned}$$

Since $a_k^2 + a_{-k}^2 = -2k_0^2 + 2e_{\mathbf{k}}^2 = -2a_k a_{-k} + 4e_{\mathbf{k}}^2$ one obtains

$$\begin{aligned}
\prod_{k_0 \in \frac{\pi}{\beta}(2\mathbf{Z}+1)}^{\epsilon} \tfrac{a_k a_{-k} + b a_k + c a_{-k} + d}{a_k a_{-k}} &= \prod_{k_0 > 0} \tfrac{(a_k a_{-k})^2 + p a_k a_{-k} + q}{(a_k a_{-k})^2} \; e^{\frac{\beta}{2}(b+c)} \\
&= \prod_{k_0 > 0} \left(1 + \tfrac{\omega_+}{a_k a_{-k}}\right) \left(1 + \tfrac{\omega_-}{a_k a_{-k}}\right) \; e^{\frac{\beta}{2}(b+c)}
\end{aligned}$$

where $p = -2e_{\mathbf{k}}(b+c) + 2d + (b+c)^2 - 4bc$ and $q = 4bc e_{\mathbf{k}}^2 - 2e_{\mathbf{k}} d(b+c) + d^2$. Application of (III.25) proves part (a) of the lemma.

Part b) First, if $b_k = a_k - \sqrt{\lambda}\, v$, one computes that

$$\mathrm{Pf} S_k = (b_k b_{-k} + \lambda \rho^2)^2 + (b_k^2 + b_{-k}^2 + 2\lambda\rho^2)\lambda(w^2 + x^2) + \lambda^2 (w^2 + x^2)^2 \tag{A.7}$$

Namely, since

$$A_{k\uparrow} A_{-k\downarrow} = a_k a_{-k} - \sqrt{\lambda}\, v(a_k + a_{-k}) + i\sqrt{\lambda}\, w(a_k - a_{-k}) + \lambda(v^2 + w^2) \quad \Rightarrow$$

$$\begin{aligned}
(A_{k\uparrow} A_{-k\downarrow} + \lambda\rho^2)(A_{k\downarrow} A_{-k\uparrow} + \lambda\rho^2) &= \Big[a_k a_{-k} - \sqrt{\lambda}\, v(a_k + a_{-k}) + \lambda(v^2 + w^2 + \rho^2)\Big]^2 \\
&\quad + \lambda w^2 (a_k - a_{-k})^2 \\
&= \big[b_k b_{-k} + \lambda(w^2 + \rho^2)\big]^2 + \lambda w^2 (b_k - b_{-k})^2
\end{aligned}$$

and

$$A_{k\uparrow} A_{k\downarrow} = (a_k - \sqrt{\lambda}\, v)^2 + \lambda w^2 = b_k^2 + \lambda w^2$$

one has

$$\begin{aligned}
\mathrm{Pf} S_k &= \big[b_k b_{-k} + \lambda(w^2 + \rho^2)\big]^2 + \lambda w^2 (b_k - b_{-k})^2 + \big[b_k^2 + b_{-k}^2 + 2\lambda(w^2 + \rho^2)\big]\lambda x^2 + \lambda^2 x^4 \\
&= \big[b_k b_{-k} + \lambda\rho^2\big]^2 + (b_k^2 + b_{-k}^2 + 2\lambda\rho^2)\lambda(w^2 + x^2) + \lambda^2 (w^2 + x^2)^2
\end{aligned}$$

Now let $y^2 = w^2 + x^2$ and

$$F_k(a_k, a_{-k}) = \big[b_k b_{-k} + \lambda\rho^2\big]^2 + (b_k^2 + b_{-k}^2 + 2\lambda\rho^2)\lambda y^2 + \lambda^2 y^4$$

Then part (b) follows from the following formula

$$\lim_{\substack{\epsilon\to 0\\ \epsilon>0}} \prod_{\substack{k_0\in\frac{\pi}{\beta}(2\mathbf{Z}+1)\\ k_0>0}} \tfrac{F_k(e^{i\epsilon k_0}a_k, e^{-i\epsilon k_0}a_{-k})}{a_k^2 a_{-k}^2} = \tfrac{\cosh^2\left(\frac{\beta}{2}\sqrt{(e_{\mathbf{k}}+\sqrt{\lambda}\,v)^2+\lambda\rho^2}\,\right)-\sin^2\left(\frac{\beta}{2}\sqrt{\lambda y^2}\right)}{\cosh^2\frac{\beta}{2}e_{\mathbf{k}}}\, e^{-\beta\sqrt{\lambda}v} \tag{A.8}$$

Proof of (A.8): One has $A_k A_{-k} = a_k a_{-k} - \sqrt{\lambda}\,v(a_k + a_{-k}) + \lambda v^2$ and

$$A_k^2 + A_{-k}^2 = a_k^2 + a_{-k}^2 - 2\sqrt{\lambda}\,v(a_k + a_{-k}) + 2\lambda v^2 = 2A_kA_{-k} + (a_k - a_{-k})^2$$

which gives

$$\begin{aligned} F_k &= (A_kA_{-k} + \lambda\rho^2 + \lambda y^2)^2 + (a_k - a_{-k})^2\lambda y^2 \\ &= [A_kA_{-k} + \lambda\rho^2 + \lambda y^2 + i\lambda^{\frac{1}{2}}y(a_k - a_{-k})]\,[A_kA_{-k} + \lambda\rho^2 + \lambda y^2 - i\lambda^{\frac{1}{2}}y(a_k - a_{-k})] \\ &= \big[a_ka_{-k} - (\sqrt{\lambda}\,v - i\lambda^{\frac{1}{2}}y)a_k - (\sqrt{\lambda}\,v + i\lambda^{\frac{1}{2}}y)a_{-k} + \lambda v^2 + \lambda\rho^2 + \lambda y^2\big]\times \\ &\quad\ \big[a_ka_{-k} - (\sqrt{\lambda}\,v + i\lambda^{\frac{1}{2}}y)a_k - (\sqrt{\lambda}\,v - i\lambda^{\frac{1}{2}}y)a_{-k} + \lambda v^2 + \lambda\rho^2 + \lambda y^2\big] \end{aligned}$$

Therefore one obtains

$$\begin{aligned} \prod_{k_0>0}^{\epsilon} \tfrac{F_k(a_k,a_{-k})}{a_k^2a_{-k}^2} &= \lim_{\substack{\epsilon\to 0\\ \epsilon<0}} \prod_{k_0} \tfrac{a_ka_{-k}-(\sqrt{\lambda}\,v-i\lambda^{\frac{1}{2}}y)e^{ik_0\epsilon}a_k-(\sqrt{\lambda}\,v+i\lambda^{\frac{1}{2}}y)e^{-ik_0\epsilon}a_{-k}+\lambda v^2+\lambda\rho^2+\lambda y^2}{a_ka_{-k}} \\ &\overset{(\mathrm{A.})}{=} \tfrac{\cosh\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\omega_+}}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\,\tfrac{\cosh\frac{\beta}{2}\sqrt{e_{\mathbf{k}}^2+\omega_-}}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\, e^{-\beta\sqrt{\lambda}\,v} \end{aligned}$$

where

$$\begin{aligned} p &= (e_{\mathbf{k}} + 2\sqrt{\lambda}\,v)^2 - e_{\mathbf{k}}^2 - 4(\lambda v^2 + \lambda y^2) + 2\lambda v^2 + 2\lambda\rho^2 + 2\lambda y^2 \\ &= 4\sqrt{\lambda}\,ve_{\mathbf{k}} + 2\lambda v^2 + 2\lambda\rho^2 - 2\lambda y^2 \\ q &= 4(\lambda v^2 + \lambda y^2)e_{\mathbf{k}}^2 + 4\sqrt{\lambda}\,ve_{\mathbf{k}}(\lambda v^2 + \lambda\rho^2 + \lambda y^2) + (\lambda v^2 + \lambda\rho^2 + \lambda y^2)^2 \\ &= (2\sqrt{\lambda}\,ve_{\mathbf{k}} + \lambda v^2 + \lambda\rho^2 + \lambda y^2)^2 + 4\lambda y^2e_{\mathbf{k}}^2 \end{aligned}$$

such that

$$\tfrac{p^2}{4} - q = -4(2\sqrt{\lambda}\,ve_{\mathbf{k}} + \lambda v^2 2 + \lambda\rho^2)\lambda y^2 - 4\lambda y^2e_{\mathbf{k}}^2 = -4\big[(e_{\mathbf{k}} + \sqrt{\lambda}\,v)^2 + \lambda\rho^2\big]\lambda y^2$$

and therefore

$$\begin{aligned} e_{\mathbf{k}}^2 + \omega_\pm &= (e_{\mathbf{k}} + \sqrt{\lambda}\,v)^2 + \lambda\rho^2 - \lambda y^2 \pm 2i\lambda^{\frac{1}{2}}y\big[(e_{\mathbf{k}} + \sqrt{\lambda}\,v)^2 + \lambda\rho^2\big]^{\frac{1}{2}} \\ &= \Big\{\big[(e_{\mathbf{k}} + \sqrt{\lambda}\,v)^2 + \lambda\rho^2\big]^{\frac{1}{2}} \pm i\lambda^{\frac{1}{2}}y\Big\}^2 \end{aligned}$$

which results in

$$\prod_{k_0>0}^{\epsilon} \frac{F_k(a_k,a_{-k})}{a_k^2 a_{-k}^2} = \frac{\cosh\frac{\beta}{2}\left\{[(e_{\mathbf{k}}+\sqrt{\lambda}\,v)^2+\lambda\rho^2]^{\frac{1}{2}}+i\lambda^{\frac{1}{2}}y\right\}}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\,\frac{\cosh\frac{\beta}{2}\left\{[(e_{\mathbf{k}}+\sqrt{\lambda}\,v)^2+\lambda\rho^2]^{\frac{1}{2}}-i\lambda^{\frac{1}{2}}y\right\}}{\cosh\frac{\beta}{2}e_{\mathbf{k}}}\,e^{-\beta\sqrt{\lambda}\,v}$$

$$= \frac{\sinh^2(\frac{\beta}{2}i\sqrt{\lambda}\,y)+\cosh^2(\frac{\beta}{2}\sqrt{(e_{\mathbf{k}}+\sqrt{\lambda}\,v)^2+\lambda\rho^2}\,)}{\cosh^2\frac{\beta}{2}e_{\mathbf{k}}}\,e^{-\beta\sqrt{\lambda}\,v} \qquad \bullet$$

References

[AB] P.W. Anderson, W.F. Brinkman, *Theory of Anisotropic Superfluidity* in *Basic Notions of Condensed Matter Physics* by P.W. Anderson, Benjamin/Cummings, 1984.

[B] N.N. Bogoliubov, *On Some Problems of the Theory of Superconductivity*, Physica 26, Supplement, p.1-16 (1960).

[BZT] N.N. Bogoliubov, D.B. Zubarev, Iu.A. Tserkovnikov, *On the Theory of Phase Transitions*, Sov. Phys. Doklady 2, p.535 (1957).

[BR] J. Bardeen, G. Rickayzen, *Ground State Energy and Green's Function for Reduced Hamiltonian for Superconductivity*, Phys. Rev. 118, p.936 (1960).

[BW] R. Balian, N.R. Werthamer, *Superconductivity with Pairs in a Relative p Wave*, Phys. Rev. 131, p.1553-1564, 1963.

[FKLT] J. Feldman, H. Knörrer, D. Lehmann, E. Trubowitz, *Fermi Liquids in Two Space Dimensions*, in *Constructive Physics*, Springer Lecture Notes in Physics, Bd. 446, 1994.

[FKT1] J. Feldman, H. Knörrer, E. Trubowitz, *Mathematical Methods of Many Body Quantum Field Theory*, Lecture Notes, ETH Zürich.

[FKT2] J. Feldman, H. Knörrer, E. Trubowitz, *A Remark on Anisotropic Superconducting States*, Helv. Phys. Acta 64, p.695-699, (1991).

[FST] J. Feldman, M. Salmhofer, E. Trubowitz, *Perturbation Theory around Non-nested Fermi Surfaces I. Keeping the Fermi Surface Fixed*, J.Stat.Phys. 84, p.1209-1336, (1996).

[FT] J. Feldman, E. Trubowitz, *Perturbation Theory for Many Fermion Systems*, Helvetia Physica Acta 63, p.156-260 (1990); *The Flow of an Electron-Phonon System to the Superconducting State*, Helv. Phys. Acta 64, p.214-357, (1991).

[G] M.D. Girardeau, *Variational Method for the Quantum Statistics of Many Particle Systems*, Phys. Rev. A 42, p.3303, (1990).

[Ha] R. Haag, *The Mathematical Structure of the Bardeen Cooper Schrieffer model*, Nuovo Cimento 25, p.287 (1962).

[Hn] E.R. Hanson, *A Table of Series and Products*, Prentice-Hall, 1975, Sec. 89.5.

[Sch] A. Schütte, *The Symmetry of the Gap in the BCS Model for Higher l-Wave Interactions*, Diploma thesis, ETH Zürich, 1997.

[W] G. Wentzel, Phys. Rev. 120, p.1572, (1960).